

Mn-doped II-VI quantum dots: artificial molecular magnets

J. Fernández-Rossier^{*1} and R. Aguado²

¹ Departamento de Física Aplicada, Universidad de Alicante, 03690 Spain

² Instituto de Ciencia de Materiales de Madrid, CSIC, 28049 Madrid, Spain

Received 1 May 2006, accepted 14 June

Published online 24 November 2006

Key words Quantum magnet, Quantum Dot, Diluted Magnetic Semiconductor

PACS 04A25

The notion of artificial atom relies on the capability to change the number of carriers one by one in semiconductor quantum dots, and the resulting changes in their electronic structure. Organic molecules with transition metal atoms that have a net magnetic moment and display hysteretic behaviour are known as single molecule magnets (SMM). The fabrication of CdTe quantum dots chemically doped with a controlled number of Mn atoms and with a number of carriers controlled either electrically or optically paves the way towards a new concept in nanomagnetism: the artificial single molecule magnet. Here we study the magnetic properties of a Mn-doped CdTe quantum dot for different charge states and show to what extent they behave like a single molecule magnet.

Copyright line will be provided by the publisher

1 Introduction

One of the most studied single molecule magnets (SMM), Mn_{12} , is made of a cage of one hundred Carbon, Hydrogen and Oxygen atoms, surrounding a $\text{Mn}_{12}\text{O}_{12}$ cluster [1]. The oxydation state of the Mn atoms either +3 or +4 so that their spin is either ($S=2$) or ($S=3/2$). In the lowest energy configuration, the molecule has total spin $S = 10$ that results from the super-exchange interactions between the Mn spins [2]. The $S = 10$ manifold has 21 states that, due to the spin-orbit interaction, are split according to $E = -DS_z^2$, where S_z is the spin projection along a crystallographic axis z [2, 3]. Remarkably, the magnetization of a molecular crystal of Mn_{12} shows hysteresis, at temperatures larger than the inter-molecular coupling, reflecting that each individual molecule behaves as a magnet [3]. In marked contrast to mono-domain magnets, the hysteresis curve of a Mn_{12} crystal displays steps at well defined values of the applied field. These are associated to quantum tunneling events of the magnetic moment. Therefore, the magnetic moment of SMM behaves quantum mechanically [4].

As an extension of the concept of artificial atom [5], we propose the notion of artificial SMM: a II-VI semiconductor quantum dot, chemically doped with Mn atoms and electrically doped with carriers so that they form a collective spin that has meta-stable states (hysteresis) and steps in the magnetization. As a substitutional impurity of the cation in the II-VI material [6], Mn is in a $2+$ oxydation state with spin $S = 5/2$. In our device there are a few Mn atoms, sufficiently apart as to make the antiferromagnetic exchange interaction between them completely negligible. The magnetic properties of the dot are given by the controlled presence of electrons and/or holes in the conduction and valence band quantum dot levels. Since both electrons and holes are exchanged coupled to the Mn spin [6], indirect exchange interactions couple the Mn spins to each other [7] in the dot. Even in the case of a dot doped with a single Mn atom, we show that the anisotropic spin interaction of the holes and the Mn [8, 9, 10, 11, 12, 13] results in SMM behaviour.

* Corresponding author: e-mail: jfrossier@ua.es, Phone: +00 999 999 999, Fax: +00 999 999 999

Most of the working principles of this proposal have been verified experimentally. On one side, the charge state of II-VI quantum dot can be prepared, with a single electron accuracy, both by electrical [14, 15, 16] and optical means [17, 18]. On the other side, several groups have reported the fabrication of II-VI quantum dots doped with Mn atoms, and the exchange coupling of electrons and holes to the Mn spins [19, 20, 21, 22, 23, 24]. More recently, the controlled fabrication of CdTe quantum dots doped with a single Mn atom [8, 9, 10], as well as the electrical control of the charge state of a single CdTe quantum dot doped with a single Mn atom [13] and a GaAs island doped with tens of Mn atoms [25] have been reported. Electrical control of the carrier density has been demonstrated in larger magnetic semiconductor structures, making it possible to alter reversibly properties of the material like the T_C [26, 27] and the coercive field [28] of these systems. Our proposal targets the achievement of this notion at the scale of a single electron and the single Mn atom limit.

Related theory work has focused on the electronic structure [7, 29, 30, 31, 32], optical response [11, 33, 34] and transport [12, 35] of Mn-doped quantum dots as a function of the number of confined carriers. It is now well established that, depending on the number of carriers present in the dot, the eigenstates of the Mn-fermion Hamiltonian are very different [7]. In previous work we have derived from numerical diagonalizations [12] the effective spin Hamiltonian for relevant low energy sector of a CdTe dot doped with a single Mn interacting with up to 3 carriers, either electrons or holes. Here we give a preliminary account of the magnetization curves of a single CdTe dot, doped with a Mn atom, as a function of the number of carriers. The results presented here permit to claim that Mn-doped quantum dots with extra holes behave like a nanomagnet, with hysteretic magnetization and steps of the magnetization at precise values of the applied field.

The rest of this paper is organized as follows. In section II we summarize the model Hamiltonian for a Mn-doped CdTe quantum dot and we review the properties of the eigenstates and eigenvalues of the Hamiltonian for a given charge state $Q = \pm 1, \pm 2, \pm 3$. In section III the magnetization of the dot with $Q = +1$ and a single Mn is calculated. We first show that the equilibrium magnetization is anisotropic, in contrast to the free Mn case, due to the interaction to the hole. Then we discuss how the magnetization behaves if the field is varied so that equilibrium can not be reached, which can be the standard situation provided the long spin relaxation time of this system.

2 Hamiltonian

We consider quantum dots with a typical size of 10 nanometers and several thousand atoms, beyond reach of present-day ab-initio calculations [31]. This justifies the use of the standard model Hamiltonian [6, 7, 11, 12, 32, 33, 34, 35] for diluted magnetic semiconductors that describes CB electrons and VB holes interacting with localized Mn spins ($\vec{M}_I$) via a local exchange interaction and their coupling to an external magnetic field, $\vec{B}$. The Mn spins interact also with each other via short range superexchange coupling. In general the Hamiltonian can not be solved exactly and in most instances mean field or some other approximations are used. Only in the limit of a few Mn atoms and a few electron and hole states considered in this paper it is possible to diagonalize the Hamiltonian numerically without approximations. The second quantization Hamiltonian of the isolated dot reads:

$$\mathcal{H}_{QD} = \sum_{\alpha, \alpha'} \left(\epsilon_{\alpha}(\vec{B}) \delta_{\alpha, \alpha'} + \sum_I J_{\alpha, \alpha'}(I) \vec{M}_I \cdot \vec{S}_{\alpha, \alpha'}(\vec{R}_I) \right) f_{\alpha}^{\dagger} f_{\alpha'} + \sum_{I, J} J_{IJ}^{\text{AF}} \vec{M}_I \cdot \vec{M}_J \quad (1)$$

Here $f_{\alpha}^{\dagger}$ creates a band carrier in the α single particle state of the quantum dot, which can be either a valence band or a conduction band state. QD carriers occupy localized spin orbitals ϕ_{α} with energy ϵ_{α} which are described in the envelope function $\vec{k} \cdot \vec{p}$ approach [7, 11, 12, 36]. In the case of valence band holes the 6 band Kohn-Luttinger Hamiltonian, including spin orbit interaction, is used as a starting point to build the quantum dot states [36]. The first term in the Hamiltonian describes non interacting carriers

in the dot and the second term describes the exchange coupling of the carriers and the Mn. We neglect interband exchange so that $J_{\alpha,\alpha'} = J_e$ ($J_{\alpha,\alpha'} = J_h$) if both α and α' belong to the conduction band states (valence band states). In contrast, we include exchange processes by which a carrier is scattered between two different levels of the dot that belong to the same band are included, although the level spacing considered here is much larger than the exchange interaction. The matrix elements of both valence and conduction band spin density, evaluated at the location of the Mn atoms, are given by $\vec{S}_{\alpha,\alpha'}(\vec{R}_I)$. They depend strongly on the orbital nature of the single particle level in question. In the case of conduction band we neglect spin orbit interactions so that $\vec{S}_{\alpha,\alpha'}$ is rotationally invariant [7]. In contrast, strong spin orbit interaction of the valence band makes the Mn-hole interaction strongly anisotropic [11, 36, 37, 38] and it varies between different dot levels. Following previous work [7, 11, 36] confinement is described by a hard wall cubic potential with $L_z < L_x \simeq L_y$. Although real dots are not cubic, this simple model [36] predicts properly that the lowest energy hole doublet has a strong heavy hole character, with spin quantized along the growth direction z whereas the next single particle doublet is mainly light hole with spin quantized in the xy plane.

As discussed in our previous work [12], in the case of $Q = \pm 1$ and $Q = \pm 3$ the low energy properties of $\mathcal{H}_{QD}$ are described by

$$\mathcal{H}_{eff} = \sum_{I,a=x,y,z} j_I^a \tau^a M_I^a + \sum_{a=x,y,z} \mu_B B_a (g M_I^a + g_a(Q) \tau^a) + \sum_{I,J} J_{IJ}^{AF} \vec{M}_I \cdot \vec{M}_J \quad (2)$$

where τ^a are the Pauli matrices operating on the isospin space defined by the lowest energy single particle doublet. In the case $Q = -1, -3$ we have $j^x = j^y = j^z$. In the case $Q = +1$ we have $j^x = j^y = 0$ in the absence of light-heavy hole mixing ($L_x = L_y$). In the case $Q = +3$ we have $j^z < j^x = j^y$.

Hereafter we only consider dots with one Mn atom. Therefore, Mn-Mn superexchange interaction is irrelevant. The coupling of Mn to conduction band electrons is spin rotational invariant, so that the ground state manifold for $Q = -1, 3$ are given by the 7 states with $S = 3$ and $S_z = \pm 3, \pm 2, \pm 1, 0$. In contrast, the Mn-hole coupling is very anisotropic. In a dot with $Q = +1$ and $L_x = L_y$ spin-flip Mn-hole interactions are strictly forbidden. The resulting Ising coupling between the Mn and the hole yields a spectrum with 6 doublets. Their wave functions are given by $|M^z, S_h^z\rangle$ with and the eigen-values $E_{M^z, S^z} = |j_z| M^z \times S^z$, where $j^z = J_h |\psi(\vec{R}_{Mn})|^2$. The lowest energy doublet corresponds to the Mn spin maximally polarized against the heavy hole spin, which lies along the growth direction, z . The addition of a second holes fills the heavy hole doublet. The low energy sector has 6 states, spin splitted according the absolute value of the third component the Mn spin, the lowest energy is given by the $M_z = \pm 1/2$ doublet. This splitting comes from inter-level exchange coupling, which are of order Δ^2/δ [7]. . The addition of a third hole into the same dot leaves a doubly occupied heavy hole doublet and a singly occupied light hole interacting with the Mn. The different spin-orbital properties of the light hole result in an exchange coupling to the Mn which is in between the isotropic Heisenberg coupling of $Q = -1$ and the Ising coupling of $Q = +1$. The picture emerging from these results is the following: the low energy states of the Mn-fermion complex are spin-split at zero applied field. The spin splitting Δ is much larger for open shell configurations than for closed shell configurations. The symmetry properties of the lowest energy states are different for electrons, heavy holes and light holes.

3 Magnetic response

Now we address the magnetic properties of the single Mn doped CdTe dot although the results also scale for larger N. We calculate the many-body spectrum of the dot as a function of an external magnetic field. Since the single particle level spacing is much larger than the cyclotron frequency, it is not a bad approximation

to neglect the orbital magnetism. The magnetization is given by :

$$\langle \vec{\mathcal{M}} \rangle = \sum_N \langle N | \rho \left(\sum_I g \mu_B \vec{M}_I + g_f \mu_B \vec{S}_f \right) | N \rangle = \langle \vec{\mathcal{M}} \rangle_{Mn} + \langle \vec{\mathcal{M}} \rangle_f \quad (3)$$

where ρ is the quantum dot density matrix and g_f denotes the gyromagnetic factor of electrons and holes.

3.1 Equilibrium Results

In a first stage we assume that the density matrix of the quantum dot is that of thermal equilibrium. We consider the case of a dot with $Q = +1$. At zero magnetic field the eigenstates with opposite magnetization are degenerate so that their populations are identical and the average magnetization vanishes. This is a consequence of the ergodic approximation, that assigns equal weight to the states of identical energy, and to time reversal symmetry. Application of a magnetic along the easy direction z splits the energy levels without mixing them, since $B_z M^z + B_z S^z$ commutes with the Hamiltonian. The resulting energy diagram is shown in figure 1, lower left panel, and features numerous level crossings, reflecting the absence of transverse interactions. The magnetic field breaks the degeneracy of all the doublets, including the lowest energy one.

At temperatures smaller than the Zeeman splitting only the ground state is occupied, resulting in a net magnetization $\langle \vec{M} \rangle$, shown in the upper panel of figure 1. We show the Mn magnetization only because is dominant and simplifies the discussion. In that figure we show the magnetization of the dot with $Q = 0$, which is the standard Brillouin function for $S = 5/2$. It is apparent that the $Q = 0$ magnetization is softer than the $Q = +1$ magnetization curve when the field is applied along the z direction, due to the exchange splitting of the levels with smaller $|M_z|$. This result is in agreement with previous work for exciton polarons [39] In contrast, the magnetization curve when the field is applied in the x direction, transverse to the easy axis of the dot, is much smaller than the free case. This is due to the competition between exchange interactions, that favor the alignment of the Mn along the z axis and the magnetic field. The difference is also seen in the energy spectrum, in the lower right panel. Hence, the presence of a heavy hole in the dot would result in a strong magnetic anisotropy in the equilibrium magnetization

3.2 Non-equilibrium results

The relaxation of the Mn-fermion spin towards equilibrium comes from the release of energy and angular momentum to their environment[40]. Since this process can be very slow, we need to consider a non-equilibrium density matrix. The relaxation time $T_1^{Q=0}$ for Mn spins in very dilute CdMnTe can be as long as 10^{-3} s[41]. In the case of a $N = Q = +1$ dot, the transitions between the two members of the ground state, $|\pm 5/2, \mp 1/2\rangle$ require a energy conserving perturbation that changes the Mn spin by $5\hbar$ and the hole angular momentum by $3\hbar$ in the opposite direction. Such a process is highly forbidden and his lifetime $T_1^{Q=+1}$ will be definitely larger than $T_1^{Q=0}$. Therefore, if the applied field is varied more rapidly than $T_1^{Q=+1}$ we are not entitled to use the equilibrium density matrix.

Now we revisit the results of the previous subsection. We assume that the Ising Hamiltonian is perturbed by very weak additional terms that will only be effective close to degeneracy points in the energy diagram of figure 2. They read:

$$\mathcal{V} = a ((M^x)^4 + (M^y)^4) + \sum_i \mathcal{J}_i \vec{M} \cdot \vec{I}_i + \epsilon (M^x \tau^x + M^y \tau^y) \quad (4)$$

The so called cubic field term is the lowest order magnetic anisotropy term allowed by the crystal symmetry [41] and makes it possible to flip the Mn spin by 4 units. The second term comes from the hyperfine coupling of the Mn d electrons to the nuclear spins $\vec{I}$ [41]. It can flip the Mn spin by one unit. The last term arises from light-hole heavy-hole mixing and permits the transition of the hole via exchange of

one unit of angular momentum with the Mn. Other terms like dipolar coupling to the nuclear spins, of phonon-mediated hole spin relaxation might be involved as well.

Our starting point is a the dot under the influence of a large magnetic field along the positive z axis at low temperatures so that we assume that the dot is in the ground state, $|+\frac{5}{2}, \downarrow\rangle$. Now the applied field is reduced down to zero, so that the state $E(+\frac{5}{2}, \downarrow)$ becomes degenerate to $E(-\frac{5}{2}, \uparrow)$. None of the terms in (4) can induce the transition from the former to the latter. It would take the combined action of several of them to achieve the spin-relaxing transition, which is thereby very slow. Hence, it would be possible to ramp the field through that degeneracy point without spin relaxation. In that case the magnetization at zero field would be finite. The field could be reversed down to the crossing point of the state $|+\frac{5}{2}, \downarrow\rangle$ with the state $|-\frac{3}{2}, \uparrow\rangle$. The cubic field term can flip the Mn spin, and it would take further spin relaxation of the hole to achieve that transition. The quantum transition could take place here, or it could happen at larger fields, where the initial state is brought in to coincidence with $|-\frac{3}{2}, \downarrow\rangle$, which is accessible via the cubic field term only. Further increase of the field towards negative value would bring into resonance the $|-\frac{3}{2}, \sigma_h\rangle$ levels with the $|-\frac{5}{2}, \uparrow\rangle$, completing the magnetization reversal process. In any case the magnetization displays remanence, hysteresis and abrupt steps as we show in the lower panel of figure 2. These are very similar to those observed in single molecule magnets [2, 3, 4]. This phenomenology is the consequence of the blocking of magnetization relaxation processes except at well defined values of the applied field that put levels with different M_z in resonance.

4 Discussion and Conclusions

Artificial atoms mimic the properties of natural atoms at much larger (smaller) length (energy) scale. From that point of view, artificial SMM is a hybrid device. The magnetic moment of both artificial and standard SMM comes from the Mn atoms. What makes them different is the glue that determines the total magnetic moment of the SMM. The properties of Mn_{12} come from the chemical bonding between the Mn and the oxygen atoms in the central core of the molecule. In the case of artificial SMM, the glue electrons are delocalized in a much larger length scale.

Measurement of the quantum features in the magnetization of molecular magnets is greatly simplified due to the homogeneity of the single molecule magnets in macroscopic molecular crystals. In the case of artificial molecular magnets, these specific values of the field that result in steps in the magnetization depend on the Mn-hole exchange coupling constant, which depends both on the dot size and on the Mn location inside the dot. Therefore, the measurement of magnetization over an ensemble of dots would yield results different from those shown in figure 3.

The measurement of the effect predicted here would require either the fabrication of extremely monodisperse dots, using for instance colloidal techniques [42], or the measurement of the magnetization of a single device. The second possibility is starting to be explored experimentally in the case of Mn_{12} molecules [43, 44]. We have studied recently the theory of a similar device in which Mn_{12} is replaced by a single Mn-doped CdTe quantum dot. We have shown that the magnetic state of the Mn determines the conductance of the dot. This opens the door for the measurement of $M(B)$ in a single dot. An additional experimental route could come from spin-selective optical measurements of a single dot. In summary, we claim that a Mn-doped CdTe quantum dot charged with a single heavy hole will have exotic magnetic properties that correspond to those of single molecule magnets. A single electron transistor with a single-Mn doped CdTe quantum dot would behave like an artificial single molecule magnet.

Acknowledgements This work has been financially supported by MEC-Spain (Grants FIS200402356, MAT2005-07369-C03-03, and the Ramon y Cajal Program) and by Generalitat Valenciana (GV05-152). This work has been partly funded by FEDER funds.




Fig. 1 Upper panel: equilibrium magnetization for dot with $Q = 0$ and $Q = +1$ with the field applied both along the growth axis and in-plane. Lower Panel: Energy spectrum for the dot as a function of B_z (left) and B_x (right).




Fig. 2 Upper panel: detail of the Mn-hole spectrum and a possible non-ergodic evolution of the density matrix. Lower panel: non-ergodic magnetization of the system.

References

- [1] T. Lis, Acta Crystallogr. Sect. B **36**, 2042 (1980)
- [2] D. Gatteschi and R. Sessoli, Angew. Chem. **42**, 268 (2003)
- [3] J. R. Friedman *et al.*, Phys. Rev. Lett. **76**, 3830 (1996)
- [4] J. Tejada, E. M. Chudnovsky, *Macroscopic Quantum Tunneling of the Magnetic Moment*, Cambridge University Press, Cambridge (1998)
- [5] P. Hawrylak, Phys. Rev. Lett. **71**, 3347 (1993) L.P. Kouwenhoven *et al.* Science **278**, 1788 (1997)
- [6] J. K. Furdyna, J. Appl. Phys. **64** R29 (1988).
- [7] J. Fernández-Rossier and L. Brey, Phys. Rev. Lett. **93** 117201 (2004)
- [8] L. Besombes, Y. Lger, L. Maingault, D. Ferrand, H. Mariette and J. Cibert Phys. Rev. Lett. **93**, 207403 (2004)
- [9] L. Besombes, Y. Leger, L. Maingault, D. Ferrand, H. Mariette, and J. Cibert Phys. Rev. B. **71**, 161307 (2005)
- [10] Y. Léger L. Besombes, L. Maingault, D. Ferrand, and H. Mariette, Phys. Rev. Lett. **95**, 047403 (2005)
- [11] J. Fernández-Rossier, Phys. Rev. B. **73**, 045301 (2006)
- [12] J. Fernández-Rossier, R. Aguado, cond-mat/0604437
- [13] Y. Léger L. Besombes, J. Fernández-Rossier, L. Maingault, H. Mariette, Phys. Rev. Lett. **97**, 107401 (2006).
- [14] D. L. Klein, R. Roth, A. K.L. Lim, A. P. Alivisatos, and P. L. McEuen Nature **389**, 699 (1997)
- [15] J. Seufert, M. Rambach, G. Bacher, A. Forchel, T. Passow, D. Hommel, Appl. Phys. Lett. **82**, 3946 (2003)
- [16] Yi Cui *et al.*, Nano Lett. **5**, 1519 (2005)
- [17] T. Flissikowski, A. Hundt, M. Lowisch, M. Rabe, and F. Henneberger Phys. Rev. Lett. **86**, 3172-3175 (2001)
- [18] L. Besombes, K. Kheng, L. Marsal, H. Mariette, Phys. Rev. B **65**, 121314 (2002)
- [19] A. A. Maksimov *et al.* Phys. Rev. B **62**, R7767 (2000)
- [20] J. Seufert *et al.*, Phys. Rev. Lett. **88** 027402, (2002)
- [21] G. Bacher *et al.*, Phys. Rev. Lett. **89**, 127201 (2002)
- [22] P. S. Dorozhkin *et al.*, Phys. Rev. B **68**, 195313 (2003)
- [23] S. Mackowski *et al.* Appl. Phys. Lett. **84**, 3337 (2004).
- [24] S. C. Erwin *et al.*, Nature **436**, 91 (2005)
- [25] J. Wunderlich *et al.* Phys. Rev. Lett. **97**, 077201 (2006)
- [26] H. Ohno *et al.*, Nature, **408**, 944 (2000)
- [27] H. Boukari *et al.*, Phys. Rev. Lett. **88**, 207204 (2002).
- [28] D. Chiba *et al.*, Science **301**, 943 (2003)
- [29] J. I. Climente, M. Korkusinski, P. Hawrylak, and J. Planelles Phys. Rev. B **71**, 125321 (2005)

- [30] F. Qu, P. Hawrylak, Phys. Rev. Lett. **95**, 217206 (2005)
- [31] X. Huang, A. Makmal, J. R. Chelikowsky, and L. Kronik, Phys. Rev. Lett. **94**, 236801 (2005)
- [32] A. O. Govorov, Phys. Rev. B **72**, 075359 (2005). A. O. Govorov, Phys. Rev. B **72**, 075358 (2005).
- [33] A. K. Bhattacharjee and J. Pérez-Conde Phys. Rev. B **68**, 045303 (2003)
- [34] A. O. Govorov, Phys. Rev. B **70**, 035321 (2004) A. O. Govorov and A. V. Kalameitsev Phys. Rev. B **71**, 035338 (2005)
- [35] A. Efros, E. Rashba, M. Rosen, Phys. Rev. Lett. **87**, 206601 (2001)
- [36] F. V. Kyrychenko and J. Kossut, Phys. Rev. B **70**, 205317 (2004)
- [37] J. Fernández-Rossier and L. J. Sham, Phys. Rev. B **64**, 235323 (2001)
- [38] L. Brey, C. Tejedor and J. Fernández-Rossier Appl. Phys. Lett. **85**, 1996 (2004)
- [39] A. K. Bhattacharjee and Phys. Rev. B **51**, 9912 (1995)
- [40] E. M. Chudnovsky and L. Gunther, Phys. Rev. Lett. **60**, 661 (1988) E. M. Chudnovsky, Phys. Rev. Lett **72**, 3433 (1994)
- [41] J. Lambe and C. Kikuchi, Phys. Rev. **119**, 1256 (1960). D. Scalbert *et al.*, Solid State Communications **66**, 571 (1988).
- [42] N. S. Norberg, D. R. Gamelin J. Phys. Chem. B **109**, 20810 (2005)
- [43] H. B. Heersche *et al.*, Phys. Rev. Lett. **96**, 206801 (2006)
- [44] Moon-Ho Jo *et al.*, cond-mat/0603276

Long title

L. First Author^{*1}, **L. Second Author**^{**1,2}, and **L. Third Author**^{***2}

¹ First address

² Second address

Received 15 November 2003, revised 30 November 2003, accepted 2 December 2003

Published online 3 December 2003

Key words List, of, comma, separated, keywords.

PACS 04A25

Copyright line will be provided by the publisher

1 First section

1.1 First subsection

Acknowledgements An acknowledgement may be placed at the end of the article.

The style of the following references should be used in all documents.

References

[1] xxx.

* Corresponding author: e-mail: x.y@xxx.yyy.zz, Phone: +00 999 999 999, Fax: +00 999 999 999

** Second author footnote.

*** Third author footnote.

Sample paper for the w-art class

L. First Author^{*1}, L. Second Author^{1,2}, and L. Third Author²

¹ First address

² Second address

Received 15 November 2003, revised 30 November 2003, accepted 2 December 2003

Published online 3 December 2003

Key words List, of, comma, separated, keywords.

PACS XXXX

This paper is a sample adapted from the $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ document testmath.tex to illustrate the use of the document class w-art and publication-specific variants of that class for WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim.

Copyright line will be provided by the publisher

1 Introduction

This paper contains examples of various features from $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

2 Enumeration of Hamiltonian paths in a graph

Let $\mathbf{A} = (a_{ij})$ be the adjacency matrix of graph G . The corresponding Kirchhoff matrix $\mathbf{K} = (k_{ij})$ is obtained from $\mathbf{A}$ by replacing in $-\mathbf{A}$ each diagonal entry by the degree of its corresponding vertex; i.e., the i th diagonal entry is identified with the degree of the i th vertex. It is well known that

$$\det \mathbf{K}(i|i) = \text{the number of spanning trees of } G, \quad i = 1, \dots, n \quad (1)$$

where $\mathbf{K}(i|i)$ is the i th principal submatrix of $\mathbf{K}$.

$\det \mathbf{K}(i|i) = \text{the number of spanning trees of } G$,

Let $C_{i(j)}$ be the set of graphs obtained from G by attaching edge $(v_i v_j)$ to each spanning tree of G . Denote by $C_i = \bigcup_j C_{i(j)}$. It is obvious that the collection of Hamiltonian cycles is a subset of C_i . Note that the cardinality of C_i is $k_{ii} \det \mathbf{K}(i|i)$. Let $\hat{X} = \{\hat{x}_1, \dots, \hat{x}_n\}$.

where $\mathbf{X} = \{\hat{x}_1, \dots, \hat{x}_n\}$

Define multiplication for the elements of $\hat{X}$ by

$$\hat{x}_i \hat{x}_j = \hat{x}_j \hat{x}_i, \quad \hat{x}_i^2 = 0, \quad i, j = 1, \dots, n. \quad (2)$$

Let $\hat{k}_{ij} = k_{ij} \hat{x}_j$ and $\hat{k}_{ij} = -\sum_{j \neq i} \hat{k}_{ij}$. Then the number of Hamiltonian cycles H_c is given by the relation [8]

$$\left(\prod_{j=1}^n \hat{x}_j \right) H_c = \frac{1}{2} \hat{k}_{ij} \det \hat{\mathbf{K}}(i|i), \quad i = 1, \dots, n. \quad (3)$$

* Corresponding author: e-mail: x.y@xxx.yyy.zz, Phone: +00 999 999 999, Fax: +00 999 999 999

The task here is to express (3) in a form free of any $\hat{x}_i, i = 1, \dots, n$. The result also leads to the resolution of enumeration of Hamiltonian paths in a graph.

It is well known that the enumeration of Hamiltonian cycles and paths in a complete graph K_n and in a complete bipartite graph $K_{n_1 n_2}$ can only be found from *first combinatorial principles* [4]. One wonders if there exists a formula which can be used very efficiently to produce K_n and $K_{n_1 n_2}$. Recently, using Lagrangian methods, Goulden and Jackson have shown that H_c can be expressed in terms of the determinant and permanent of the adjacency matrix [3]. However, the formula of Goulden and Jackson determines neither K_n nor $K_{n_1 n_2}$ effectively. In this paper, using an algebraic method, we parametrize the adjacency matrix. The resulting formula also involves the determinant and permanent, but it can easily be applied to K_n and $K_{n_1 n_2}$. In addition, we eliminate the permanent from H_c and show that H_c can be represented by a determinantal function of multivariables, each variable with domain $\{0, 1\}$. Furthermore, we show that H_c can be written by number of spanning trees of subgraphs. Finally, we apply the formulas to a complete multigraph $K_{n_1 \dots n_p}$.

The conditions $a_{ij} = a_{ji}, i, j = 1, \dots, n$, are not required in this paper. All formulas can be extended to a digraph simply by multiplying H_c by 2.

3 Main Theorem

Notation For $p, q \in P$ and $n \in \omega$ we write $(q, n) \leq (p, n)$ if $q \leq p$ and $A_{q,n} = A_{p,n}$.

\begin{notation} For $p, q \in P$ and $n \in \omega$

...

\end{notation}

Let $\mathbf{B} = (b_{ij})$ be an $n \times n$ matrix. Let $\mathbf{n} = \{1, \dots, n\}$. Using the properties of (2), it is readily seen that

Lemma 3.1

$$\prod_{i \in \mathbf{n}} \left(\sum_{j \in \mathbf{n}} b_{ij} \hat{x}_i \right) = \left(\prod_{i \in \mathbf{n}} \hat{x}_i \right) \text{per } \mathbf{B} \quad (4)$$

where $\text{per } \mathbf{B}$ is the permanent of $\mathbf{B}$.

Let $\hat{Y} = \{\hat{y}_1, \dots, \hat{y}_n\}$. Define multiplication for the elements of $\hat{Y}$ by

$$\hat{y}_i \hat{y}_j + \hat{y}_j \hat{y}_i = 0, \quad i, j = 1, \dots, n. \quad (5)$$

Then, it follows that

Lemma 3.2

$$\prod_{i \in \mathbf{n}} \left(\sum_{j \in \mathbf{n}} b_{ij} \hat{y}_j \right) = \left(\prod_{i \in \mathbf{n}} \hat{y}_i \right) \det \mathbf{B}. \quad (6)$$

Note that all basic properties of determinants are direct consequences of Lemma 3.2. Write

$$\sum_{j \in \mathbf{n}} b_{ij} \hat{y}_j = \sum_{j \in \mathbf{n}} b_{ij}^{(\lambda)} \hat{y}_j + (b_{ii} - \lambda_i) \hat{y}_i \quad (7)$$

where

$$b_{ii}^{(\lambda)} = \lambda_i, \quad b_{ij}^{(\lambda)} = b_{ij}, \quad i \neq j. \quad (8)$$

Let $\mathbf{B}^{(\lambda)} = (b_{ij}^{(\lambda)})$. By (6) and (7), it is straightforward to show the following result:

Theorem 3.3

$$\det \mathbf{B} = \sum_{l=0}^n \sum_{I_l \subseteq \mathbf{n}} \prod_{i \in I_l} (b_{ii} - \lambda_i) \det \mathbf{B}^{(\lambda)}(I_l | I_l), \quad (9)$$

where $I_l = \{i_1, \dots, i_l\}$ and $\mathbf{B}^{(\lambda)}(I_l | I_l)$ is the principal submatrix obtained from $\mathbf{B}^{(\lambda)}$ by deleting its $i_1, \dots, i_l$ rows and columns.

Remark 3.4 Let $\mathbf{M}$ be an $n \times n$ matrix. The convention $\mathbf{M}(\mathbf{n} | \mathbf{n}) = 1$ has been used in (9) and hereafter.

Before proceeding with our discussion, we pause to note that Theorem 3.3 yields immediately a fundamental formula which can be used to compute the coefficients of a characteristic polynomial [9]:

Corollary 3.5 Write $\det(\mathbf{B} - x\mathbf{I}) = \sum_{l=0}^n (-1)^l b_l x^l$. Then

$$b_l = \sum_{I_l \subseteq \mathbf{n}} \det \mathbf{B}(I_l | I_l). \quad (10)$$

Let

$$\mathbf{K}(t, t_1, \dots, t_n) = \begin{pmatrix} D_1 t & -a_{12} t_2 & \dots & -a_{1n} t_n \\ -a_{21} t_1 & D_2 t & \dots & -a_{2n} t_n \\ \dots & \dots & \dots & \dots \\ -a_{n1} t_1 & -a_{n2} t_2 & \dots & D_n t \end{pmatrix}, \quad (11)$$

$$\begin{pmatrix} D_1 t & -a_{12} t_2 & \dots & -a_{1n} t_n \\ -a_{21} t_1 & D_2 t & \dots & -a_{2n} t_n \\ \dots & \dots & \dots & \dots \\ -a_{n1} t_1 & -a_{n2} t_2 & \dots & D_n t \end{pmatrix}$$

where

$$D_i = \sum_{j \in \mathbf{n}} a_{ij} t_j, \quad i = 1, \dots, n. \quad (12)$$

Set

$$D(t_1, \dots, t_n) = \frac{\delta}{\delta t} \det \mathbf{K}(t, t_1, \dots, t_n) \big|_{t=1}.$$

Then

$$D(t_1, \dots, t_n) = \sum_{i \in \mathbf{n}} D_i \det \mathbf{K}(t = 1, t_1, \dots, t_n; i | i), \quad (13)$$

where $\mathbf{K}(t = 1, t_1, \dots, t_n; i | i)$ is the i th principal submatrix of $\mathbf{K}(t = 1, t_1, \dots, t_n)$.

Theorem 3.3 leads to

$$\det \mathbf{K}(t_1, t_1, \dots, t_n) = \sum_{I \in \mathbf{n}} (-1)^{|I|} t^{n-|I|} \prod_{i \in I} t_i \prod_{j \in I} (D_j + \lambda_j t_j) \det \mathbf{A}^{(\lambda)}(\bar{I} | \bar{I}). \quad (14)$$

Note that

$$\det \mathbf{K}(t = 1, t_1, \dots, t_n) = \sum_{I \in \mathbf{n}} (-1)^{|I|} \prod_{i \in I} t_i \prod_{j \in I} (D_j + \lambda_j t_j) \det \mathbf{A}^{(\lambda)}(\bar{I} | \bar{I}) = 0. \quad (15)$$

Let $t_i = \hat{x}_i, i = 1, \dots, n$. Lemma 3.1 yields

$$\begin{aligned} \left(\sum_{i \in \mathbf{n}} a_{l_i} x_i \right) \det \mathbf{K}(t = 1, x_1, \dots, x_n; l|l) \\ = \left(\prod_{i \in \mathbf{n}} \hat{x}_i \right) \sum_{I \subseteq \mathbf{n} - \{l\}} (-1)^{|I|} \text{per } \mathbf{A}^{(\lambda)}(I|I) \det \mathbf{A}^{(\lambda)}(\overline{I} \cup \{l\} | \overline{I} \cup \{l\}). \end{aligned} \quad (16)$$

```
\begin{multline}
\biggl(\sum_{i \in \mathbf{n}} a_{l_i} x_i \biggr)
\det \mathbf{K}(t=1, x_1, \dots, x_n; l | l) \\
= \biggl(\prod_{i \in \mathbf{n}} \hat{x}_i \biggr)
\sum_{I \subseteq \mathbf{n} - \{l\}}
(-1)^{|\overline{I}|} \text{per } \mathbf{A}^{(\lambda)}(I|I)
\det \mathbf{A}^{(\lambda)}(\overline{I} \cup \{l\} | \overline{I} \cup \{l\}).
\label{sum-ali}
\end{multline}
```

By (3), (6), and (7), we have

Proposition 3.6

$$H_c = \frac{1}{2n} \sum_{l=0}^n (-1)^l D_l, \quad (17)$$

where

$$D_l = \sum_{I_l \subseteq \mathbf{n}} D(t_1, \dots, t_n) 2^{|I_l|} \Big|_{t_i = \begin{cases} 0, & \text{if } i \in I_l \\ 1, & \text{otherwise} \end{cases}}, \quad i = 1, \dots, n. \quad (18)$$

4 Application

We consider here the applications of Theorems 5.2 and 5.3 to a complete multipartite graph $K_{n_1 \dots n_p}$. It can be shown that the number of spanning trees of $K_{n_1 \dots n_p}$ may be written

$$T = n^{p-2} \prod_{i=1}^p (n - n_i)^{n_i-1} \quad (19)$$

where

$$n = n_1 + \dots + n_p. \quad (20)$$

It follows from Theorems 5.2 and 5.3 that

$$\begin{aligned} H_c = \frac{1}{2n} \sum_{l=0}^n (-1)^l (n-l)^{p-2} \sum_{l_1 + \dots + l_p = l} \prod_{i=1}^p \binom{n_i}{l_i} \\ \cdot [(n-l) - (n_i - l_i)]^{n_i - l_i} \cdot \left[(n-l)^2 - \sum_{j=1}^p (n_j - l_j)^2 \right]. \end{aligned} \quad (21)$$

... \binom{n_i}{l_i} \dots

and

$$H_c = \frac{1}{2} \sum_{l=0}^{n-1} (-1)^l (n-l)^{p-2} \sum_{l_1+\dots+l_p=l} \prod_{i=1}^p \binom{n_i}{l_i} \cdot [(n-l) - (n_i - l_i)]^{n_i-l_i} \left(1 - \frac{l_p}{n_p}\right) [(n-l) - (n_p - l_p)]. \quad (22)$$

The enumeration of H_c in a $K_{n_1 \dots n_p}$ graph can also be carried out by Theorem 7.5 or 7.6 together with the algebraic method of (2). Some elegant representations may be obtained. For example, H_c in a $K_{n_1 n_2 n_3}$ graph may be written

$$H_c = \frac{n_1! n_2! n_3!}{n_1 + n_2 + n_3} \sum_i \left[\binom{n_1}{i} \binom{n_2}{n_3 - n_1 + i} \binom{n_3}{n_3 - n_2 + i} + \binom{n_1 - 1}{i} \binom{n_2 - 1}{n_3 - n_1 + i} \binom{n_3 - 1}{n_3 - n_2 + i} \right]. \quad (23)$$

5 Secret Key Exchanges

Modern cryptography is fundamentally concerned with the problem of secure private communication. A Secret Key Exchange is a protocol where Alice and Bob, having no secret information in common to start, are able to agree on a common secret key, conversing over a public channel. The notion of a Secret Key Exchange protocol was first introduced in the seminal paper of Diffie and Hellman [1]. [1] presented a concrete implementation of a Secret Key Exchange protocol, dependent on a specific assumption (a variant on the discrete log), specially tailored to yield Secret Key Exchange. Secret Key Exchange is of course trivial if trapdoor permutations exist. However, there is no known implementation based on a weaker general assumption.

The concept of an informationally one-way function was introduced in [5]. We give only an informal definition here:

Definition 5.1 A polynomial time computable function $f = \{f_k\}$ is informationally one-way if there is no probabilistic polynomial time algorithm which (with probability of the form $1 - k^{-e}$ for some $e > 0$) returns on input $y \in \{0, 1\}^k$ a random element of $f^{-1}(y)$.

In the non-uniform setting [5] show that these are not weaker than one-way functions:

Theorem 5.2 ([5] (non-uniform)) *The existence of informationally one-way functions implies the existence of one-way functions.*

We will stick to the convention introduced above of saying “non-uniform” before the theorem statement when the theorem makes use of non-uniformity. It should be understood that if nothing is said then the result holds for both the uniform and the non-uniform models.

It now follows from Theorem 5.2 that

Theorem 5.3 (non-uniform) *Weak SKE implies the existence of a one-way function.*

More recently, the polynomial-time, interior point algorithms for linear programming have been extended to the case of convex quadratic programs [11, 13], certain linear complementarity problems [7, 10], and the nonlinear complementarity problem [6]. The connection between these algorithms and the classical Newton method for nonlinear equations is well explained in [7].

6 Review

We begin our discussion with the following definition:

Definition 6.1 A function $H: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be *B-differentiable* at the point z if (i) H is Lipschitz continuous in a neighborhood of z , and (ii) there exists a positive homogeneous function $BH(z): \mathbb{R}^n \rightarrow \mathbb{R}^n$, called the *B-derivative* of H at z , such that

$$\lim_{v \rightarrow 0} \frac{H(z+v) - H(z) - BH(z)v}{\|v\|} = 0.$$

The function H is *B-differentiable in set S* if it is B-differentiable at every point in S . The B-derivative $BH(z)$ is said to be *strong* if

$$\lim_{(v,v') \rightarrow (0,0)} \frac{H(z+v) - H(z+v') - BH(z)(v-v')}{\|v-v'\|} = 0.$$

Lemma 6.2 *There exists a smooth function $\psi_0(z)$ defined for $|z| > 1 - 2a$ satisfying the following properties:*

- (i) $\psi_0(z)$ is bounded above and below by positive constants $c_1 \leq \psi_0(z) \leq c_2$.
- (ii) If $|z| > 1$, then $\psi_0(z) = 1$.
- (iii) For all z in the domain of ψ_0 , $\Delta_0 \ln \psi_0 \geq 0$.
- (iv) If $1 - 2a < |z| < 1 - a$, then $\Delta_0 \ln \psi_0 \geq c_3 > 0$.

Proof. We choose $\psi_0(z)$ to be a radial function depending only on $r = |z|$. Let $h(r) \geq 0$ be a suitable smooth function satisfying $h(r) \geq c_3$ for $1 - 2a < |z| < 1 - a$, and $h(r) = 0$ for $|z| > 1 - \frac{a}{2}$. The radial Laplacian

$$\Delta_0 \ln \psi_0(r) = \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) \ln \psi_0(r)$$

has smooth coefficients for $r > 1 - 2a$. Therefore, we may apply the existence and uniqueness theory for ordinary differential equations. Simply let $\ln \psi_0(r)$ be the solution of the differential equation

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) \ln \psi_0(r) = h(r)$$

with initial conditions given by $\ln \psi_0(1) = 0$ and $\ln \psi'_0(1) = 0$.

Next, let D_ν be a finite collection of pairwise disjoint disks, all of which are contained in the unit disk centered at the origin in C . We assume that $D_\nu = \{z \mid |z - z_\nu| < \delta\}$. Suppose that $D_\nu(a)$ denotes the smaller concentric disk $D_\nu(a) = \{z \mid |z - z_\nu| \leq (1 - 2a)\delta\}$. We define a smooth weight function $\Phi_0(z)$ for $z \in C - \bigcup_\nu D_\nu(a)$ by setting $\Phi_0(z) = 1$ when $z \notin \bigcup_\nu D_\nu$ and $\Phi_0(z) = \psi_0((z - z_\nu)/\delta)$ when z is an element of D_ν . It follows from Lemma 6.2 that Φ_0 satisfies the properties:

- (i) $\Phi_0(z)$ is bounded above and below by positive constants $c_1 \leq \Phi_0(z) \leq c_2$.
- (ii) $\Delta_0 \ln \Phi_0 \geq 0$ for all $z \in C - \bigcup_\nu D_\nu(a)$, the domain where the function Φ_0 is defined.
- (iii) $\Delta_0 \ln \Phi_0 \geq c_3 \delta^{-2}$ when $(1 - 2a)\delta < |z - z_\nu| < (1 - a)\delta$.

Let A_ν denote the annulus $A_\nu = \{(1 - 2a)\delta < |z - z_\nu| < (1 - a)\delta\}$, and set $A = \bigcup_\nu A_\nu$. The properties (2) and (3) of Φ_0 may be summarized as $\Delta_0 \ln \Phi_0 \geq c_3 \delta^{-2} \chi_A$, where χ_A is the characteristic function of A . $\square$

Suppose that α is a nonnegative real constant. We apply Proposition 3.6 with $\Phi(z) = \Phi_0(z)e^{\alpha|z|^2}$. If $u \in C_0^\infty(R^2 - \bigcup_\nu D_\nu(a))$, assume that $\mathcal{D}$ is a bounded domain containing the support of u and $A \subset \mathcal{D} \subset R^2 - \bigcup_\nu D_\nu(a)$. A calculation gives

$$\int_{\mathcal{D}} |\bar{\partial}u|^2 \Phi_0(z) e^{\alpha|z|^2} \geq c_4 \alpha \int_{\mathcal{D}} |u|^2 \Phi_0 e^{\alpha|z|^2} + c_5 \delta^{-2} \int_A |u|^2 \Phi_0 e^{\alpha|z|^2}.$$

The boundedness, property (1) of Φ_0 , then yields

$$\int_{\mathcal{D}} |\bar{\partial}u|^2 e^{\alpha|z|^2} \geq c_6 \alpha \int_{\mathcal{D}} |u|^2 e^{\alpha|z|^2} + c_7 \delta^{-2} \int_A |u|^2 e^{\alpha|z|^2}.$$

Let $B(X)$ be the set of blocks of Λ_X and let $b(X) = |B(X)|$. If $\phi \in Q_X$ then ϕ is constant on the blocks of Λ_X .

$$P_X = \{\phi \in M \mid \Lambda_\phi = \Lambda_X\}, \quad Q_X = \{\phi \in M \mid \Lambda_\phi \geq \Lambda_X\}. \quad (24)$$

If $\Lambda_\phi \geq \Lambda_X$ then $\Lambda_\phi = \Lambda_Y$ for some $Y \geq X$ so that

$$Q_X = \bigcup_{Y \geq X} P_Y.$$

Thus by Möbius inversion

$$|P_Y| = \sum_{X \geq Y} \mu(Y, X) |Q_X|.$$

Thus there is a bijection from Q_X to $W^{B(X)}$. In particular $|Q_X| = w^{b(X)}$.

Next note that $b(X) = \dim X$. We see this by choosing a basis for X consisting of vectors v^k defined by

$$v_i^k = \begin{cases} 1 & \text{if } i \in \Lambda_k, \\ 0 & \text{otherwise.} \end{cases}$$

$\backslash [v^{\wedge \{k\}}_{\{i\}} =$
 $\backslash \begin{cases} 1 & \text{if } i \in \Lambda_{\{k\}} \\ 0 & \text{otherwise.} \end{cases} \backslash \backslash$
 $\backslash]$

Lemma 6.3 *Let $\mathcal{A}$ be an arrangement. Then*

$$\chi(\mathcal{A}, t) = \sum_{\mathcal{B} \subseteq \mathcal{A}} (-1)^{|\mathcal{B}|} t^{\dim T(\mathcal{B})}.$$

In order to compute R'' recall the definition of $S(X, Y)$ from Lemma 3.1. Since $H \in \mathcal{B}$, $\mathcal{A}_H \subseteq \mathcal{B}$. Thus if $T(\mathcal{B}) = Y$ then $\mathcal{B} \in S(H, Y)$. Let $L'' = L(\mathcal{A}'')$. Then

$$\begin{aligned} R'' &= \sum_{H \in \mathcal{B} \subseteq \mathcal{A}} (-1)^{|\mathcal{B}|} t^{\dim T(\mathcal{B})} \\ &= \sum_{Y \in L''} \sum_{\mathcal{B} \in S(H, Y)} (-1)^{|\mathcal{B}|} t^{\dim Y} \\ &= - \sum_{Y \in L''} \sum_{\mathcal{B} \in S(H, Y)} (-1)^{|\mathcal{B} - \mathcal{A}_H|} t^{\dim Y} \\ &= - \sum_{Y \in L''} \mu(H, Y) t^{\dim Y} \\ &= -\chi(\mathcal{A}'', t). \end{aligned} \quad (25)$$

Corollary 6.4 *Let $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ be a triple of arrangements. Then*

$$\pi(\mathcal{A}, t) = \pi(\mathcal{A}', t) + t\pi(\mathcal{A}'', t).$$

Definition 6.5 *Let $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ be a triple with respect to the hyperplane $H \in \mathcal{A}$. Call H a *separator* if $T(\mathcal{A}) \notin L(\mathcal{A}')$.*

Corollary 6.6 *Let $(\mathcal{A}, \mathcal{A}', \mathcal{A}'')$ be a triple with respect to $H \in \mathcal{A}$.*

(i) *If H is a separator then*

$$\mu(\mathcal{A}) = -\mu(\mathcal{A}'')$$

and hence

$$|\mu(\mathcal{A})| = |\mu(\mathcal{A}'')|.$$

(ii) *If H is not a separator then*

$$\mu(\mathcal{A}) = \mu(\mathcal{A}') - \mu(\mathcal{A}'')$$

and

$$|\mu(\mathcal{A})| = |\mu(\mathcal{A}')| + |\mu(\mathcal{A}'')|.$$

Proof. It follows from Theorem 5.2 that $\pi(\mathcal{A}, t)$ has leading term

$$(-1)^{r(\mathcal{A})} \mu(\mathcal{A}) t^{r(\mathcal{A})}.$$

The conclusion follows by comparing coefficients of the leading terms on both sides of the equation in Corollary 6.4. If H is a separator then $r(\mathcal{A}') < r(\mathcal{A})$ and there is no contribution from $\pi(\mathcal{A}', t)$. $\square$

The Poincaré polynomial of an arrangement will appear repeatedly in these notes. It will be shown to equal the Poincaré polynomial of the graded algebras which we are going to associate with $\mathcal{A}$. It is also the Poincaré polynomial of the complement $M(\mathcal{A})$ for a complex arrangement. Here we prove that the Poincaré polynomial is the chamber counting function for a real arrangement. The complement $M(\mathcal{A})$ is a disjoint union of chambers

$$M(\mathcal{A}) = \bigcup_{C \in \text{Cham}(\mathcal{A})} C.$$

The number of chambers is determined by the Poincaré polynomial as follows.

Theorem 6.7 *Let $\mathcal{A}_{\mathbf{R}}$ be a real arrangement. Then*

$$|\text{Cham}(\mathcal{A}_{\mathbf{R}})| = \pi(\mathcal{A}_{\mathbf{R}}, 1).$$

Proof. We check the properties required in Corollary 6.6: (i) follows from $\pi(\Phi_l, t) = 1$, and (ii) is a consequence of Corollary 3.5. $\square$

Theorem 6.8 *Let ϕ be a protocol for a random pair (X, Y) . If one of $\sigma_\phi(x', y)$ and $\sigma_\phi(x, y')$ is a prefix of the other and $(x, y) \in S_{X, Y}$, then*

$$\langle \sigma_j(x', y) \rangle_{j=1}^\infty = \langle \sigma_j(x, y) \rangle_{j=1}^\infty = \langle \sigma_j(x, y') \rangle_{j=1}^\infty.$$

Fig. 1 $Q(\mathcal{A}_1) = xyz(x-z)(x+z)(y-z)(y+z)$ **Fig. 2** $Q(\mathcal{A}_2) = xyz(x+y+z)(x+y-z)(x-y+z)(x-y-z)$

Proof. We show by induction on i that

$$\langle \sigma_j(x', y) \rangle_{j=1}^i = \langle \sigma_j(x, y) \rangle_{j=1}^i = \langle \sigma_j(x, y') \rangle_{j=1}^i.$$

The induction hypothesis holds vacuously for $i = 0$. Assume it holds for $i-1$, in particular $[\sigma_j(x', y)]_{j=1}^{i-1} = [\sigma_j(x, y')]_{j=1}^{i-1}$. Then one of $[\sigma_j(x', y)]_{j=i}^\infty$ and $[\sigma_j(x, y')]_{j=i}^\infty$ is a prefix of the other which implies that one of $\sigma_i(x', y)$ and $\sigma_i(x, y')$ is a prefix of the other. If the i th message is transmitted by P_X then, by the separate-transmissions property and the induction hypothesis, $\sigma_i(x, y) = \sigma_i(x, y')$, hence one of $\sigma_i(x, y)$ and $\sigma_i(x', y)$ is a prefix of the other. By the implicit-termination property, neither $\sigma_i(x, y)$ nor $\sigma_i(x', y)$ can be a proper prefix of the other, hence they must be the same and $\sigma_i(x', y) = \sigma_i(x, y) = \sigma_i(x, y')$. If the i th message is transmitted by P_Y then, symmetrically, $\sigma_i(x, y) = \sigma_i(x', y)$ by the induction hypothesis and the separate-transmissions property, and, then, $\sigma_i(x, y) = \sigma_i(x, y')$ by the implicit-termination property, proving the induction step. $\square$

If ϕ is a protocol for (X, Y) , and $(x, y), (x', y)$ are distinct inputs in $S_{X,Y}$, then, by the correct-decision property, $\langle \sigma_j(x, y) \rangle_{j=1}^\infty \neq \langle \sigma_j(x', y) \rangle_{j=1}^\infty$.

Equation (25) defined P_Y 's ambiguity set $S_{X|Y}(y)$ to be the set of possible X values when $Y = y$. The last corollary implies that for all $y \in S_Y$, the multiset¹ of codewords $\{\sigma_\phi(x, y) : x \in S_{X|Y}(y)\}$ is prefix free.

¹ A multiset allows multiplicity of elements. Hence, $\{0, 01, 01\}$ is prefix free as a set, but not as a multiset.

7 One-Way Complexity

$\hat{C}_1(X|Y)$, the one-way complexity of a random pair (X, Y) , is the number of bits P_X must transmit in the worst case when P_Y is not permitted to transmit any feedback messages. Starting with $S_{X,Y}$, the support set of (X, Y) , we define $G(X|Y)$, the *characteristic hypergraph* of (X, Y) , and show that

$$\hat{C}_1(X|Y) = \lceil \log \chi(G(X|Y)) \rceil.$$

Let (X, Y) be a random pair. For each y in S_Y , the support set of Y , Equation (25) defined $S_{X|Y}(y)$ to be the set of possible x values when $Y = y$. The *characteristic hypergraph* $G(X|Y)$ of (X, Y) has S_X as its vertex set and the hyperedge $S_{X|Y}(y)$ for each $y \in S_Y$.

We can now prove a continuity theorem.

Theorem 7.1 *Let $\Omega \subset \mathbf{R}^n$ be an open set, let $u \in BV(\Omega; \mathbf{R}^m)$, and let*

$$T_x^u = \left\{ y \in \mathbf{R}^m : y = \tilde{u}(x) + \left\langle \frac{Du}{|Du|}(x), z \right\rangle \text{ for some } z \in \mathbf{R}^n \right\} \quad (26)$$

for every $x \in \Omega \setminus S_u$. Let $f: \mathbf{R}^m \rightarrow \mathbf{R}^k$ be a Lipschitz continuous function such that $f(0) = 0$, and let $v = f(u): \Omega \rightarrow \mathbf{R}^k$. Then $v \in BV(\Omega; \mathbf{R}^k)$ and

$$Jv = (f(u^+) - f(u^-)) \otimes \nu_u \cdot \mathcal{H}_{n-1}|_{S_u}. \quad (27)$$

In addition, for $|\tilde{Du}|$ -almost every $x \in \Omega$ the restriction of the function f to T_x^u is differentiable at $\tilde{u}(x)$ and

$$\tilde{D}v = \nabla(f|_{T_x^u})(\tilde{u}) \frac{\tilde{Du}}{|\tilde{Du}|} \cdot |\tilde{Du}|. \quad (28)$$

Before proving the theorem, we state without proof three elementary remarks which will be useful in the sequel.

Remark 7.2 Let $\omega:]0, +\infty[\rightarrow]0, +\infty[$ be a continuous function such that $\omega(t) \rightarrow 0$ as $t \rightarrow 0$. Then

$$\lim_{h \rightarrow 0^+} g(\omega(h)) = L \Leftrightarrow \lim_{h \rightarrow 0^+} g(h) = L$$

for any function $g:]0, +\infty[\rightarrow \mathbf{R}$.

Remark 7.3 Let $g: \mathbf{R}^n \rightarrow \mathbf{R}$ be a Lipschitz continuous function and assume that

$$L(z) = \lim_{h \rightarrow 0^+} \frac{g(hz) - g(0)}{h}$$

exists for every $z \in \mathbf{Q}^n$ and that L is a linear function of z . Then g is differentiable at 0.

Remark 7.4 Let $A: \mathbf{R}^n \rightarrow \mathbf{R}^m$ be a linear function, and let $f: \mathbf{R}^m \rightarrow \mathbf{R}$ be a function. Then the restriction of f to the range of A is differentiable at 0 if and only if $f(A): \mathbf{R}^n \rightarrow \mathbf{R}$ is differentiable at 0 and

$$\nabla(f|_{\text{Im}(A)})(0)A = \nabla(f(A))(0).$$

Proof. We begin by showing that $v \in BV(\Omega; \mathbf{R}^k)$ and

$$|Dv|(B) \leq K |Du|(B) \quad \forall B \in \mathbf{B}(\Omega), \quad (29)$$

where $K > 0$ is the Lipschitz constant of f . By (13) and by the approximation result quoted in §3, it is possible to find a sequence $(u_h) \subset C^1(\Omega; \mathbf{R}^m)$ converging to u in $L^1(\Omega; \mathbf{R}^m)$ and such that

$$\lim_{h \rightarrow +\infty} \int_{\Omega} |\nabla u_h| \, dx = |Du|(\Omega).$$

The functions $v_h = f(u_h)$ are locally Lipschitz continuous in Ω , and the definition of differential implies that $|\nabla v_h| \leq K |\nabla u_h|$ almost everywhere in Ω . The lower semicontinuity of the total variation and (13) yield

$$\begin{aligned} |Dv|(\Omega) &\leq \liminf_{h \rightarrow +\infty} |Dv_h|(\Omega) = \liminf_{h \rightarrow +\infty} \int_{\Omega} |\nabla v_h| \, dx \\ &\leq K \liminf_{h \rightarrow +\infty} \int_{\Omega} |\nabla u_h| \, dx = K |Du|(\Omega). \end{aligned} \quad (30)$$

Since $f(0) = 0$, we have also

$$\int_{\Omega} |v| \, dx \leq K \int_{\Omega} |u| \, dx;$$

therefore $u \in BV(\Omega; \mathbf{R}^k)$. Repeating the same argument for every open set $A \subset \Omega$, we get (29) for every $B \in \mathbf{B}(\Omega)$, because $|Dv|, |Du|$ are Radon measures. To prove Lemma 6.2, first we observe that

$$S_v \subset S_u, \quad \tilde{v}(x) = f(\tilde{u}(x)) \quad \forall x \in \Omega \setminus S_u. \quad (31)$$

In fact, for every $\varepsilon > 0$ we have

$$\{y \in B_{\rho}(x) : |v(y) - f(\tilde{u}(x))| > \varepsilon\} \subset \{y \in B_{\rho}(x) : |u(y) - \tilde{u}(x)| > \varepsilon/K\},$$

hence

$$\lim_{\rho \rightarrow 0^+} \frac{|\{y \in B_{\rho}(x) : |v(y) - f(\tilde{u}(x))| > \varepsilon\}|}{\rho^n} = 0$$

whenever $x \in \Omega \setminus S_u$. By a similar argument, if $x \in S_u$ is a point such that there exists a triplet (u^+, u^-, ν_u) satisfying (14), (15), then

$$(v^+(x) - v^-(x)) \otimes \nu_v = (f(u^+(x)) - f(u^-(x))) \otimes \nu_u \quad \text{if } x \in S_v$$

and $f(u^-(x)) = f(u^+(x))$ if $x \in S_u \setminus S_v$. Hence, by (1.8) we get

$$\begin{aligned} Jv(B) &= \int_{B \cap S_v} (v^+ - v^-) \otimes \nu_v \, d\mathcal{H}_{n-1} = \int_{B \cap S_v} (f(u^+) - f(u^-)) \otimes \nu_u \, d\mathcal{H}_{n-1} \\ &= \int_{B \cap S_u} (f(u^+) - f(u^-)) \otimes \nu_u \, d\mathcal{H}_{n-1} \end{aligned}$$

and Lemma 6.2 is proved. $\square$

To prove (31), it is not restrictive to assume that $k = 1$. Moreover, to simplify our notation, from now on we shall assume that $\Omega = \mathbf{R}^n$. The proof of (31) is divided into two steps. In the first step we prove the statement in the one-dimensional case ($n = 1$), using Theorem 5.3. In the second step we achieve the general result using Theorem 7.1.

Step 1

Assume that $n = 1$. Since S_u is at most countable, (7) yields that $|\tilde{D}v|(S_u \setminus S_v) = 0$, so that (19) and (21) imply that $Dv = \tilde{D}v + Jv$ is the Radon-Nikodým decomposition of Dv in absolutely continuous and singular part with respect to $|\tilde{D}u|$. By Theorem 5.3, we have

$$\frac{\tilde{D}v}{|\tilde{D}u|}(t) = \lim_{s \rightarrow t^+} \frac{Dv([t, s])}{|\tilde{D}u|([t, s])}, \quad \frac{\tilde{D}u}{|\tilde{D}u|}(t) = \lim_{s \rightarrow t^+} \frac{Du([t, s])}{|\tilde{D}u|([t, s])}$$

$|\tilde{D}u|$ -almost everywhere in $\mathbf{R}$. It is well known (see, for instance, [12, 2.5.16]) that every one-dimensional function of bounded variation w has a unique left continuous representative, i.e., a function $\hat{w}$ such that $\hat{w} = w$ almost everywhere and $\lim_{s \rightarrow t^-} \hat{w}(s) = \hat{w}(t)$ for every $t \in \mathbf{R}$. These conditions imply

$$\hat{u}(t) = Du(-\infty, t], \quad \hat{v}(t) = Dv(-\infty, t] \quad \forall t \in \mathbf{R} \quad (32)$$

and

$$\hat{v}(t) = f(\hat{u}(t)) \quad \forall t \in \mathbf{R}. \quad (33)$$

Let $t \in \mathbf{R}$ be such that $|\tilde{D}u|([t, s]) > 0$ for every $s > t$ and assume that the limits in (22) exist. By (23) and (24) we get

$$\begin{aligned} \frac{\hat{v}(s) - \hat{v}(t)}{|\tilde{D}u|([t, s])} &= \frac{f(\hat{u}(s)) - f(\hat{u}(t))}{|\tilde{D}u|([t, s])} \\ &= \frac{f(\hat{u}(s)) - f(\hat{u}(t) + \frac{\tilde{D}u}{|\tilde{D}u|}(t) |\tilde{D}u|([t, s]))}{|\tilde{D}u|([t, s])} \\ &\quad + \frac{f(\hat{u}(t) + \frac{\tilde{D}u}{|\tilde{D}u|}(t) |\tilde{D}u|([t, s])) - f(\hat{u}(t))}{|\tilde{D}u|([t, s])} \end{aligned}$$

for every $s > t$. Using the Lipschitz condition on f we find

$$\begin{aligned} \left| \frac{\hat{v}(s) - \hat{v}(t)}{|\tilde{D}u|([t, s])} - \frac{f(\hat{u}(t) + \frac{\tilde{D}u}{|\tilde{D}u|}(t) |\tilde{D}u|([t, s])) - f(\hat{u}(t))}{|\tilde{D}u|([t, s])} \right| \\ \leq K \left| \frac{\hat{u}(s) - \hat{u}(t)}{|\tilde{D}u|([t, s])} - \frac{\tilde{D}u}{|\tilde{D}u|}(t) \right|. \end{aligned}$$

By (29), the function $s \rightarrow \left| \tilde{D}u \right| ([t, s])$ is continuous and converges to 0 as $s \downarrow t$. Therefore Remark 7.2 and the previous inequality imply

$$\frac{\tilde{D}v}{\left| \tilde{D}u \right|}(t) = \lim_{h \rightarrow 0^+} \frac{f(\hat{u}(t) + h \frac{\tilde{D}u}{\left| \tilde{D}u \right|}(t)) - f(\hat{u}(t))}{h} \quad \left| \tilde{D}u \right| \text{-a.e. in } \mathbf{R}.$$

By (22), $\hat{u}(x) = \tilde{u}(x)$ for every $x \in \mathbf{R} \setminus S_u$; moreover, applying the same argument to the functions $u'(t) = u(-t)$, $v'(t) = f(u'(t)) = v(-t)$, we get

$$\frac{\tilde{D}v}{\left| \tilde{D}u \right|}(t) = \lim_{h \rightarrow 0} \frac{f(\tilde{u}(t) + h \frac{\tilde{D}u}{\left| \tilde{D}u \right|}(t)) - f(\tilde{u}(t))}{h} \quad \left| \tilde{D}u \right| \text{-a.e. in } \mathbf{R}$$

and our statement is proved.

Step 2

Let us consider now the general case $n > 1$. Let $\nu \in \mathbf{R}^n$ be such that $|\nu| = 1$, and let $\pi_\nu = \{y \in \mathbf{R}^n : \langle y, \nu \rangle = 0\}$. In the following, we shall identify $\mathbf{R}^n$ with $\pi_\nu \times \mathbf{R}$, and we shall denote by y the variable ranging in π_ν and by t the variable ranging in $\mathbf{R}$. By the just proven one-dimensional result, and by Theorem 3.3, we get

$$\lim_{h \rightarrow 0} \frac{f(\tilde{u}(y + t\nu) + h \frac{\tilde{D}u_y}{\left| \tilde{D}u_y \right|}(t)) - f(\tilde{u}(y + t\nu))}{h} = \frac{\tilde{D}v_y}{\left| \tilde{D}u_y \right|}(t) \quad \left| \tilde{D}u_y \right| \text{-a.e. in } \mathbf{R}$$

for $\mathcal{H}_{n-1}$ -almost every $y \in \pi_\nu$. We claim that

$$\frac{\langle \tilde{D}u, \nu \rangle}{\left| \langle \tilde{D}u, \nu \rangle \right|}(y + t\nu) = \frac{\tilde{D}u_y}{\left| \tilde{D}u_y \right|}(t) \quad \left| \tilde{D}u_y \right| \text{-a.e. in } \mathbf{R} \quad (34)$$

for $\mathcal{H}_{n-1}$ -almost every $y \in \pi_\nu$. In fact, by (16) and (18) we get

$$\begin{aligned} \int_{\pi_\nu} \frac{\tilde{D}u_y}{\left| \tilde{D}u_y \right|} \cdot \left| \tilde{D}u_y \right| d\mathcal{H}_{n-1}(y) &= \int_{\pi_\nu} \tilde{D}u_y d\mathcal{H}_{n-1}(y) \\ &= \langle \tilde{D}u, \nu \rangle = \frac{\langle \tilde{D}u, \nu \rangle}{\left| \langle \tilde{D}u, \nu \rangle \right|} \cdot \left| \langle \tilde{D}u, \nu \rangle \right| = \int_{\pi_\nu} \frac{\langle \tilde{D}u, \nu \rangle}{\left| \langle \tilde{D}u, \nu \rangle \right|}(y + \cdot \nu) \cdot \left| \tilde{D}u_y \right| d\mathcal{H}_{n-1}(y) \end{aligned}$$

and (24) follows from (13). By the same argument it is possible to prove that

$$\frac{\langle \tilde{D}v, \nu \rangle}{\left| \langle \tilde{D}u, \nu \rangle \right|}(y + t\nu) = \frac{\tilde{D}v_y}{\left| \tilde{D}u_y \right|}(t) \quad \left| \tilde{D}u_y \right| \text{-a.e. in } \mathbf{R} \quad (35)$$

for $\mathcal{H}_{n-1}$ -almost every $y \in \pi_\nu$. By (24) and (25) we get

$$\lim_{h \rightarrow 0} \frac{f(\tilde{u}(y + t\nu) + h \frac{\langle \tilde{D}u, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(y + t\nu)) - f(\tilde{u}(y + t\nu))}{h} = \frac{\langle \tilde{D}v, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(y + t\nu)$$

for $\mathcal{H}_{n-1}$ -almost every $y \in \pi_\nu$, and using again (14), (15) we get

$$\lim_{h \rightarrow 0} \frac{f(\tilde{u}(x) + h \frac{\langle \tilde{D}u, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(x)) - f(\tilde{u}(x))}{h} = \frac{\langle \tilde{D}v, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(x)$$

$|\langle \tilde{D}u, \nu \rangle|$ -a.e. in $\mathbf{R}^n$.

Since the function $|\langle \tilde{D}u, \nu \rangle| / |\tilde{D}u|$ is strictly positive $|\langle \tilde{D}u, \nu \rangle|$ -almost everywhere, we obtain also

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(\tilde{u}(x) + h \frac{|\langle \tilde{D}u, \nu \rangle|}{|\tilde{D}u|}(x) \frac{\langle \tilde{D}u, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(x)) - f(\tilde{u}(x))}{h} \\ = \frac{|\langle \tilde{D}u, \nu \rangle|}{|\tilde{D}u|}(x) \frac{\langle \tilde{D}v, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|}(x) \end{aligned}$$

$|\langle \tilde{D}u, \nu \rangle|$ -almost everywhere in $\mathbf{R}^n$.

Finally, since

$$\begin{aligned} \frac{|\langle \tilde{D}u, \nu \rangle|}{|\tilde{D}u|} \frac{\langle \tilde{D}u, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|} &= \frac{\langle \tilde{D}u, \nu \rangle}{|\tilde{D}u|} = \left\langle \frac{\tilde{D}u}{|\tilde{D}u|}, \nu \right\rangle \quad |\tilde{D}u| \text{-a.e. in } \mathbf{R}^n \\ \frac{|\langle \tilde{D}u, \nu \rangle|}{|\tilde{D}u|} \frac{\langle \tilde{D}v, \nu \rangle}{|\langle \tilde{D}u, \nu \rangle|} &= \frac{\langle \tilde{D}v, \nu \rangle}{|\tilde{D}u|} = \left\langle \frac{\tilde{D}v}{|\tilde{D}u|}, \nu \right\rangle \quad |\tilde{D}u| \text{-a.e. in } \mathbf{R}^n \end{aligned}$$

and since both sides of (33) are zero $|\tilde{D}u|$ -almost everywhere on $|\langle \tilde{D}u, \nu \rangle|$ -negligible sets, we conclude that

$$\lim_{h \rightarrow 0} \frac{f\left(\tilde{u}(x) + h \left\langle \frac{\tilde{D}u}{|\tilde{D}u|}(x), \nu \right\rangle\right) - f(\tilde{u}(x))}{h} = \left\langle \frac{\tilde{D}v}{|\tilde{D}u|}(x), \nu \right\rangle,$$

$|\tilde{D}u|$ -a.e. in $\mathbf{R}^n$. Since ν is arbitrary, by Remarks 7.3 and 7.4 the restriction of f to the affine space T_x^u is differentiable at $\tilde{u}(x)$ for $|\tilde{D}u|$ -almost every $x \in \mathbf{R}^n$ and (26) holds. $\square$

It follows from (13), (14), and (15) that

$$D(t_1, \dots, t_n) = \sum_{I \in \mathbf{n}} (-1)^{|I|-1} |I| \prod_{i \in I} t_i \prod_{j \in I} (D_j + \lambda_j t_j) \det \mathbf{A}^{(\lambda)}(\bar{I}|\bar{I}). \quad (36)$$

Let $t_i = \hat{x}_i$, $i = 1, \dots, n$. Lemma 1 leads to

$$D(\hat{x}_1, \dots, \hat{x}_n) = \prod_{i \in \mathbf{n}} \hat{x}_i \sum_{I \in \mathbf{n}} (-1)^{|I|-1} |I| \text{per } \mathbf{A}^{(\lambda)}(I|I) \det \mathbf{A}^{(\lambda)}(\bar{I}|\bar{I}). \quad (37)$$

By (3), (13), and (37), we have the following result:

Theorem 7.5

$$H_c = \frac{1}{2n} \sum_{l=1}^n l (-1)^{l-1} A_l^{(\lambda)}, \quad (38)$$

where

$$A_l^{(\lambda)} = \sum_{I_l \subseteq \mathbf{n}} \text{per } \mathbf{A}^{(\lambda)}(I_l|I_l) \det \mathbf{A}^{(\lambda)}(\bar{I}_l|\bar{I}_l), |I_l| = l. \quad (39)$$

It is worth noting that $A_l^{(\lambda)}$ of (39) is similar to the coefficients b_l of the characteristic polynomial of (10). It is well known in graph theory that the coefficients b_l can be expressed as a sum over certain subgraphs. It is interesting to see whether A_l , $\lambda = 0$, structural properties of a graph.

We may call (38) a parametric representation of H_c . In computation, the parameter λ_i plays very important roles. The choice of the parameter usually depends on the properties of the given graph. For a complete graph K_n , let $\lambda_i = 1$, $i = 1, \dots, n$. It follows from (39) that

$$A_l^{(1)} = \begin{cases} n!, & \text{if } l = 1 \\ 0, & \text{otherwise.} \end{cases} \quad (40)$$

By (38)

$$H_c = \frac{1}{2} (n-1)!. \quad (41)$$

For a complete bipartite graph $K_{n_1 n_2}$, let $\lambda_i = 0$, $i = 1, \dots, n$. By (39),

$$A_l = \begin{cases} -n_1! n_2! \delta_{n_1 n_2}, & \text{if } l = 2 \\ 0, & \text{otherwise.} \end{cases} \quad (42)$$

Theorem 7.5 leads to

$$H_c = \frac{1}{n_1 + n_2} n_1! n_2! \delta_{n_1 n_2}. \quad (43)$$

Now, we consider an asymmetrical approach. Theorem 3.3 leads to

$$\begin{aligned} \det \mathbf{K}(t = 1, t_1, \dots, t_n; l|l) \\ = \sum_{I \subseteq \mathbf{n} - \{l\}} (-1)^{|I|} \prod_{i \in I} t_i \prod_{j \in I} (D_j + \lambda_j t_j) \det \mathbf{A}^{(\lambda)}(\bar{I} \cup \{l\} | \bar{I} \cup \{l\}). \end{aligned} \quad (44)$$

By (3) and (16) we have the following asymmetrical result:

Theorem 7.6

$$H_c = \frac{1}{2} \sum_{I \subseteq \mathbf{n} - \{l\}} (-1)^{|I|} \text{per } \mathbf{A}^{(\lambda)}(I|I) \det \mathbf{A}^{(\lambda)}(\bar{I} \cup \{l\} | \bar{I} \cup \{l\}) \quad (45)$$

which reduces to Goulden–Jackson’s formula when $\lambda_i = 0$, $i = 1, \dots, n$ [9].

8 Various font features of the `amsmath` package

8.1 Bold versions of special symbols

In the `amsmath` package `\boldsymbol` is used for getting individual bold math symbols and bold Greek letters—everything in math except for letters of the Latin alphabet, where you'd use `\mathbf`. For example,

```
A_\infty + \pi A_0 \sim
\mathbf{A}_\{\boldsymbol{\infty}\} \boldsymbol{+}
\boldsymbol{\pi} \mathbf{A}_\{\boldsymbol{0}\}
```

looks like this:

$$A_\infty + \pi A_0 \sim \mathbf{A}_\infty + \pi \mathbf{A}_0$$

8.2 “Poor man’s bold”

If a bold version of a particular symbol doesn’t exist in the available fonts, then `\boldsymbol` can’t be used to make that symbol bold. At the present time, this means that `\boldsymbol` can’t be used with symbols from the `msam` and `msbm` fonts, among others. In some cases, poor man’s bold (`\pmb`) can be used instead of `\boldsymbol`:

$$\frac{\partial x}{\partial y} \bigg| \frac{\partial y}{\partial z}$$

```
\[\frac{\partial x}{\partial y} \bigg| \frac{\partial y}{\partial z}\]
\pmb{\bigg\vert}
\frac{\partial y}{\partial z}
```

So-called “large operator” symbols such as $\sum$ and $\prod$ require an additional command, `\mathop`, to produce proper spacing and limits when `\pmb` is used. For further details see *The T_EXbook*.

$$\sum_{\substack{i < B \\ i \text{ odd}}} \prod_{\kappa} \kappa F(r_i) \quad \sum_{\substack{i < B \\ i \text{ odd}}} \prod_{\kappa} \kappa(r_i)$$

```
\[\sum_{\substack{i < B \\ i \text{ odd}}} \prod_{\kappa} \kappa F(r_i) \quad
\mathop{\pmb{\sum}}_{\substack{i < B \\ i \text{ odd}}} \prod_{\kappa} \kappa(r_i)
\]
```

9 Compound symbols and other features

9.1 Multiple integral signs

`\iint`, `\iiint`, and `\iiint` give multiple integral signs with the spacing between them nicely adjusted, in both text and display style. `\idotsint` gives two integral signs with dots between them.

$$\int_A \int f(x, y) dx dy \quad \int_A \int \int f(x, y, z) dx dy dz \quad (46)$$

$$\int_A \int \int \int f(w, x, y, z) dw dx dy dz \quad \int_A \cdots \int f(x_1, \dots, x_k) \quad (47)$$

9.2 Over and under arrows

Some extra over and under arrow operations are provided in the `amsmath` package. (Basic $\text{\LaTeX}$ provides `\overrightarrow` and `\overleftarrow`).

$$\begin{aligned}\overrightarrow{\psi_\delta(t)E_th} &= \overrightarrow{\psi_\delta(t)E_th} \\ \overleftarrow{\psi_\delta(t)E_th} &= \overleftarrow{\psi_\delta(t)E_th} \\ \overleftrightarrow{\psi_\delta(t)E_th} &= \overleftrightarrow{\psi_\delta(t)E_th}\end{aligned}$$

```
\begin{align*}
\overrightarrow{\psi_\delta(t) E_t h}&
=\underrightarrow{\psi_\delta(t) E_t h}\backslash
\overleftarrow{\psi_\delta(t) E_t h}&
=\underleftarrow{\psi_\delta(t) E_t h}\backslash
\overleftrightharpoonup{\psi_\delta(t) E_t h}&
=\underleftrightharpoonup{\psi_\delta(t) E_t h}
\end{align*}
```

These all scale properly in subscript sizes:

$$\int_{\overrightarrow{AB}} ax \, dx$$

```
\[\int_{\overrightarrow{AB}} ax\,dx\]
```

9.3 Dots

Normally you need only type `\dots` for ellipsis dots in a math formula. The main exception is when the dots fall at the end of the formula; then you need to specify one of `\dotsc` (series dots, after a comma), `\dotsb` (binary dots, for binary relations or operators), `\dotsm` (multiplication dots), or `\dotsi` (dots after an integral). For example, the input

Then we have the series `$A_1,A_2,\dotsc$`,
the regional sum `$A_1+A_2+\dotsb$`,
the orthogonal product `$A_1A_2\dotsm$`,
and the infinite integral
`\[\int_{A_1}\int_{A_2}\dotsi\]`.

produces

Then we have the series $A_1, A_2, \dots$, the regional sum $A_1 + A_2 + \dots$, the orthogonal product $A_1 A_2 \dots$, and the infinite integral

$$\int_{A_1} \int_{A_2} \dots$$

9.4 Accents in math

Double accents:

$$\hat{H} \quad \check{C} \quad \tilde{T} \quad \acute{A} \quad \grave{G} \quad \dot{D} \quad \ddot{D} \quad \breve{B} \quad \bar{B} \quad \vec{V}$$

$$\begin{array}{l} \backslash \backslash \text{Hat}\{\text{Hat}\{H\}\} \backslash \text{quad} \backslash \text{Check}\{\backslash \text{Check}\{C\}\} \backslash \text{quad} \\ \backslash \text{Tilde}\{\backslash \text{Tilde}\{T\}\} \backslash \text{quad} \backslash \text{Acute}\{\backslash \text{Acute}\{A\}\} \backslash \text{quad} \\ \backslash \text{Grave}\{\backslash \text{Grave}\{G\}\} \backslash \text{quad} \backslash \text{Dot}\{\backslash \text{Dot}\{D\}\} \backslash \text{quad} \\ \backslash \text{Ddot}\{\backslash \text{Ddot}\{D\}\} \backslash \text{quad} \backslash \text{Breve}\{\backslash \text{Breve}\{B\}\} \backslash \text{quad} \\ \backslash \text{Bar}\{\backslash \text{Bar}\{B\}\} \backslash \text{quad} \backslash \text{Vec}\{\backslash \text{Vec}\{V\}\} \backslash \backslash \end{array}$$

This double accent operation is complicated and tends to slow down the processing of a $\text{\LaTeX}$ file.

9.5 Dot accents

`\dddot` and `\ddddot` are available to produce triple and quadruple dot accents in addition to the `\dot` and `\ddot` accents already available in L^AT_EX:

$$\ddot{Q} \quad \ddot{R}$$
$$\llbracket \text{ddd}\dot{\{Q\}} \rrbracket \text{qquad} \llbracket \text{ddd}\dot{\{R\}} \rrbracket$$

9.6 Roots

In the `amsmath` package `\leftroot` and `\uproot` allow you to adjust the position of the root index of a radical:

$$\sqrt[\leftroot{-2}\uproot{2}\beta]{k}$$

gives good positioning of the β :

 $\sqrt[k]{\beta}$

9.7 Boxed formulas

The command `\boxed` puts a box around its argument, like `\fbox` except that the contents are in math mode:

$$\boxed{W_t - F \subseteq V(P_i) \subseteq W_t}$$

$$\boxed{W_t - F \subseteq V(P_i) \subseteq W_t}.$$

9.8 Extensible arrows

`\leftarrow` and `\rightarrow` produce arrows that extend automatically to accommodate unusually wide subscripts or superscripts. The text of the subscript or superscript are given as an optional resp. mandatory argument: Example:

$$0 \xleftarrow[\zeta]{\alpha} F \times \Delta[n-1] \xrightarrow{\partial_0 \alpha(b)} E^{\partial_0 b}$$

$$\left[0 \rightarrow \Omega^{\alpha} F \rightarrow \Omega^{\alpha} E \rightarrow \Omega^{\alpha} \mathcal{O}_B \rightarrow 0\right]$$

9.9 `\overset`, `\underset`, and `\sideset`

Examples:

$$\overset{*}{X} \quad \underset{*}{X} \quad \overset{a}{\underset{b}{X}}$$

```
\[\overset{*}{X}\qquad\underset{*}{X}\qquad
\overset{a}{\underset{b}{X}}\]
```

The command `\sideset` is for a rather special purpose: putting symbols at the subscript and superscript corners of a large operator symbol such as $\sum$ or $\prod$, without affecting the placement of limits. Examples:

$$\prod_k^* \sum_{0 \leq i \leq m}' E_i \beta x$$

```
\[\sideset{_{*}^{*}}{\prod_k}\qquad
\sideset{}{'}\sum_{0\leq i\leq m} E_i\beta x
\]
```

9.10 The `\text` command

The main use of the command `\text` is for words or phrases in a display:

$$\mathbf{y} = \mathbf{y}' \quad \text{if and only if} \quad y'_k = \delta_k y_{\tau(k)}$$

```
\[\mathbf{y}=\mathbf{y}'\quad\text{if and only if}\quad
y'_k=\delta_k y_{\tau(k)}\]
```

9.11 Operator names

The more common math functions such as $\log$, $\sin$, and $\lim$ have predefined control sequences: `\log`, `\sin`, `\lim`. The `amsmath` package provides `\DeclareMathOperator` and `\DeclareMathOperator*` for producing new function names that will have the same typographical treatment. Examples:

$$\|f\|_{\infty} = \operatorname{ess\,sup}_{x \in R^n} |f(x)|$$

```
\[\norm{f}_{\infty}=
\operatorname{esssup}_{x\in R^n}\abs{f(x)}\]
```

$$\operatorname{meas}_1\{u \in R_+^1 : f^*(u) > \alpha\} = \operatorname{meas}_n\{x \in R^n : |f(x)| \geq \alpha\} \quad \forall \alpha > 0.$$

```
\[\operatorname{meas}_1\{u\in R_+^1\colon f^*(u)>\alpha\}
=\operatorname{meas}_n\{x\in R^n\colon \abs{f(x)}\geq\alpha\}
\quad\forall\alpha>0.\]
```

`\esssup` and `\meas` would be defined in the document preamble as

```
\DeclareMathOperator*\esssup{ess\, sup}
\DeclareMathOperator{\meas}{meas}
```

The following special operator names are predefined in the `amsmath` package: `\varlimsup`, `\varliminf`, `\varinjlim`, and `\varprojlim`. Here's what they look like in use:

$$\overline{\lim}_{n \rightarrow \infty} \mathcal{Q}(u_n, u_n - u^\#) \leq 0 \quad (48)$$

$$\varliminf_{n \rightarrow \infty} |a_{n+1}| / |a_n| = 0 \quad (49)$$

$$\varinjlim (m_i^\lambda \cdot)^* \leq 0 \quad (50)$$

$$\varprojlim_{p \in S(A)} A_p \leq 0 \quad (51)$$

```
\begin{align}
&\&\varlimsup_{n \rightarrow \infty} \mathcal{Q}(u_n, u_n - u^\#) \leq 0 \\
&\&\varliminf_{n \rightarrow \infty} |a_{n+1}| / |a_n| = 0 \\
&\&\varinjlim (m_i^\lambda \cdot)^* \leq 0 \\
&\&\varprojlim_{p \in S(A)} A_p \leq 0 \\
\end{align}
```

9.12 `\mod` and its relatives

The commands `\mod` and `\pod` are variants of `\pmod` preferred by some authors; `\mod` omits the parentheses, whereas `\pod` omits the ‘mod’ and retains the parentheses. Examples:

$$x \equiv y + 1 \pmod{m^2} \quad (52)$$

$$x \equiv y + 1 \mod m^2 \quad (53)$$

$$x \equiv y + 1 \pod{m^2} \quad (54)$$

```
\begin{align}
x &\equiv y+1 \pmod{m^2} \\
x &\equiv y+1 \mod{m^2} \\
x &\equiv y+1 \pod{m^2} \\
\end{align}
```

9.13 Fractions and related constructions

The usual notation for binomials is similar to the fraction concept, so it has a similar command `\binom` with two arguments. Example:

$$\begin{aligned} \sum_{\gamma \in \Gamma_C} I_\gamma &= 2^k - \binom{k}{1} 2^{k-1} + \binom{k}{2} 2^{k-2} \\ &\quad + \cdots + (-1)^l \binom{k}{l} 2^{k-l} + \cdots + (-1)^k \\ &= (2-1)^k = 1 \end{aligned} \quad (55)$$

```
\begin{equation}
\begin{split}
&[\sum_{\gamma \in \Gamma_C} I_\gamma \\
&= 2^k - \binom{k}{1} 2^{k-1} + \binom{k}{2} 2^{k-2} \\
&\quad + \cdots + (-1)^l \binom{k}{l} 2^{k-l} + \cdots + (-1)^k
\end{split}
\end{equation}
```

```
+ \dots + (-1)^k \\
&= (2-1)^k = 1 \\
\end{split} \\
\end{equation}
```

There are also abbreviations

```
\dfrac          \dbinom
\tfrac          \tbinom
```

for the commonly needed constructions

```
{\displaystyle\frac ... }   {\displaystyle\binom ... }
{\textstyle\frac ... }      {\textstyle\binom ... }
```

The generalized fraction command `\genfrac` provides full access to the six TeX fraction primitives:

$$\backslash\over: \frac{n+1}{2} \qquad \backslash\overwithdelims: \left\langle \frac{n+1}{2} \right\rangle \quad (56)$$

$$\backslash\atop: \frac{n+1}{2} \qquad \backslash\atopwithdelims: \binom{n+1}{2} \quad (57)$$

$$\backslash\above: \frac{n+1}{2} \qquad \backslash\abovewithdelims: \left[\frac{n+1}{2} \right] \quad (58)$$

```
\text{\cn{over}: } & \backslash\genfrac{}{}{}{}{n+1}{2} &
\text{\cn{overwithdelims}: } &
\backslash\genfrac{\langle}{\rangle}{}{}{n+1}{2} \\
\text{\cn{atop}: } & \backslash\genfrac{}{}{0pt}{}{n+1}{2} &
\text{\cn{atopwithdelims}: } &
\backslash\genfrac{({)}{}{0pt}{}{n+1}{2} \\
\text{\cn{above}: } & \backslash\genfrac{}{}{1pt}{}{n+1}{2} &
\text{\cn{abovewithdelims}: } &
\backslash\genfrac{[]{}{1pt}{}{n+1}{2}
```

9.14 Continued fractions

The continued fraction

$$\frac{1}{\sqrt{2} + \frac{1}{\sqrt{2} + \frac{1}{\sqrt{2} + \frac{1}{\sqrt{2} + \frac{1}{\sqrt{2} + \dots}}}}} \quad (59)$$

can be obtained by typing

```
\cfrac{1}{\sqrt{2}+
\cfrac{1}{\sqrt{2}+
\cfrac{1}{\sqrt{2}+
\cfrac{1}{\sqrt{2}+
\cfrac{1}{\sqrt{2}+\dotsb
}}}}}
```

Left or right placement of any of the numerators is accomplished by using `\cfrac[l]` or `\cfrac[r]` instead of `\cfrac`.

9.15 Smash

In `amsmath` there are optional arguments `t` and `b` for the plain `TEX` command `\smash`, because sometimes it is advantageous to be able to ‘smash’ only the top or only the bottom of something while retaining the natural depth or height. In the formula $X_j = (1/\sqrt{\lambda_j})X'_j$ `\smash[b]` has been used to limit the size of the radical symbol.

`$X_j=(1/\sqrt{\smash[b]{\lambda_j}})X'_j$`

Without the use of `\smash[b]` the formula would have appeared thus: $X_j = (1/\sqrt{\lambda_j})X'_j$, with the radical extending to encompass the depth of the subscript j .

9.16 The ‘cases’ environment

‘Cases’ constructions like the following can be produced using the `CASES` environment.

$$P_{r-j} = \begin{cases} 0 & \text{if } r-j \text{ is odd,} \\ r!(-1)^{(r-j)/2} & \text{if } r-j \text{ is even.} \end{cases} \quad (60)$$

```
\begin{equation} P_{r-j}=
\begin{cases}
0& \text{\text{if } $r-j$ is odd},\\
r!\,(-1)^{(r-j)/2}& \text{\text{if } $r-j$ is even}.
\end{cases}
\end{equation}
```

Notice the use of `\text` and the embedded math.

9.17 Matrix

Here are samples of the matrix environments, `\matrix`, `\pmatrix`, `\bmatrix`, `\Bmatrix`, `\vmatrix` and `\Vmatrix`:

$$\vartheta \quad \varrho \quad \begin{pmatrix} \vartheta & \varrho \end{pmatrix} \quad \begin{bmatrix} \vartheta & \varrho \end{bmatrix} \quad \left\{ \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right\} \quad \left| \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right| \quad \left\| \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right\| \quad (61)$$

```
\begin{matrix}
\vartheta& \varrho& \begin{pmatrix} \vartheta & \varrho \end{pmatrix} & \begin{bmatrix} \vartheta & \varrho \end{bmatrix} & \left\{ \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right\} & \left| \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right| & \left\| \begin{matrix} \vartheta & \varrho \\ \varphi & \varpi \end{matrix} \right\|
\end{matrix}
```

To produce a small matrix suitable for use in text, use the `smallmatrix` environment.

```
\begin{math}
  \bigl( \begin{smallmatrix}
    a&b\\
    c&d
  \end{smallmatrix} \bigr)
\end{math}
```

To show the effect of the matrix on the surrounding lines of a paragraph, we put it here: $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and follow it with enough text to ensure that there will be at least one full line below the matrix.

`\hdotsfor{number}` produces a row of dots in a matrix spanning the given number of columns:

$$W(\Phi) = \begin{vmatrix} \frac{\varphi}{(\varphi_1, \varepsilon_1)} & 0 & \dots & 0 \\ \frac{\varphi^{k_{n2}}}{(\varphi_2, \varepsilon_1)} & \frac{\varphi}{(\varphi_2, \varepsilon_2)} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ \frac{\varphi^{k_{n1}}}{(\varphi_n, \varepsilon_1)} & \frac{\varphi^{k_{n2}}}{(\varphi_n, \varepsilon_2)} & \dots & \frac{\varphi^{k_{nn-1}}}{(\varphi_n, \varepsilon_{n-1})} & \frac{\varphi}{(\varphi_n, \varepsilon_n)} \end{vmatrix}$$

```
\[W(\Phi)= \begin{Vmatrix}
\dfrac{\varphi_{\varphi_1,\varepsilon_1}}{(\varphi_1,\varepsilon_1)}&\dots&0\\
\dfrac{\varphi_{k_{n2}}}{(\varphi_2,\varepsilon_1)}&\frac{\varphi}{(\varphi_2,\varepsilon_2)}&\dots&0\\
\dotsfor{5}\\
\dfrac{\varphi_{k_{n1}}}{(\varphi_n,\varepsilon_1)}&\frac{\varphi^{k_{n2}}}{(\varphi_n,\varepsilon_2)}&\dots&\frac{\varphi^{k_{nn-1}}}{(\varphi_n,\varepsilon_{n-1})}&\frac{\varphi}{(\varphi_n,\varepsilon_n)}
\end{Vmatrix}\]
```

The spacing of the dots can be varied through use of a square-bracket option, for example, `\hdotsfor[1.5]{3}`. The number in square brackets will be used as a multiplier; the normal value is 1.

9.18 The `\substack` command

The `\substack` command can be used to produce a multiline subscript or superscript: for example

```
\sum_{\substack{0\leq i\leq m\\ 0<j<n}} P(i,j)
```

produces a two-line subscript underneath the sum:

$$\sum_{\substack{0\leq i\leq m\\ 0<j<n}} P(i,j) \quad (62)$$

A slightly more generalized form is the `subarray` environment which allows you to specify that each line should be left-aligned instead of centered, as here:

$$\sum_{\subarray{0\leq i\leq m\\ 0<j<n}} P(i,j) \quad (63)$$

```
\sum_{\begin{subarray}{l}
  0\leq i\leq m\\
  0<j<n
\end{subarray}}
P(i,j)
```

9.19 Big-g-g delimiters

Here are some big delimiters, first in `\normalsize`:

$$\left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds\right)$$

```
\[\biggl(\mathbf{E}_{\{y\}}
\int_0^{t_{\varepsilon}} L_{\{x,y^x(s)\}} \varphi(x) \, ds
\biggr)
\]
```

and now in `\Large` size:

$$\left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds\right)$$

```
{\Large
\[\biggl(\mathbf{E}_{\{y\}}
\int_0^{t_{\varepsilon}} L_{\{x,y^x(s)\}} \varphi(x) \, ds
\biggr)
\]}
```


A Examples of multiple-line equation structures

Note: Starting on this page, vertical rules are added at the margins so that the positioning of various display elements with respect to the margins can be seen more clearly.

A.1 Split

The `split` environment is not an independent environment but should be used inside something else such as `equation` or `align`.

If there is not enough room for it, the equation number for a `split` will be shifted to the previous line, when equation numbers are on the left; the number shifts down to the next line when numbers are on the right.

$$\begin{aligned}
 f_{h,\varepsilon}(x,y) &= \varepsilon \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon u)} \varphi(x) du \\
 &= h \int L_{x,z} \varphi(x) \rho_x(dz) \\
 &\quad + h \left[\frac{1}{t_\varepsilon} \left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds - t_\varepsilon \int L_{x,z} \varphi(x) \rho_x(dz) \right) \right. \\
 &\quad \left. + \frac{1}{t_\varepsilon} \left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds - \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon s)} \varphi(x) ds \right) \right] \\
 &= h \widehat{L}_x \varphi(x) + h \theta_\varepsilon(x,y),
 \end{aligned} \tag{64}$$

Some text after to test the below-display spacing.

```

\begin{equation}
\begin{split}
f_{h,\varepsilon}(x,y)
&= \varepsilon \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon u)} \varphi(x) du \\
&= h \int L_{x,z} \varphi(x) \rho_x(dz) \\
&\quad + h \biggl[ \frac{1}{t_\varepsilon} \biggl( \mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds \\
&\quad - t_\varepsilon \int L_{x,z} \varphi(x) \rho_x(dz) \biggr) \\
&\quad + \frac{1}{t_\varepsilon} \biggl( \mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds \\
&\quad - \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon s)} \varphi(x) ds \biggr) \biggr] \\
&= h \widehat{L}_x \varphi(x) + h \theta_\varepsilon(x,y),
\end{split}
\end{equation}

```

Unnumbered version:

$$\begin{aligned}
 f_{h,\varepsilon}(x,y) &= \varepsilon \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon u)} \varphi(x) du \\
 &= h \int L_{x,z} \varphi(x) \rho_x(dz) \\
 &\quad + h \left[\frac{1}{t_\varepsilon} \left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds - t_\varepsilon \int L_{x,z} \varphi(x) \rho_x(dz) \right) \right. \\
 &\quad \left. + \frac{1}{t_\varepsilon} \left(\mathbf{E}_y \int_0^{t_\varepsilon} L_{x,y^x(s)} \varphi(x) ds - \mathbf{E}_{x,y} \int_0^{t_\varepsilon} L_{x,y_\varepsilon(\varepsilon s)} \varphi(x) ds \right) \right] \\
 &= h \hat{L}_x \varphi(x) + h \theta_\varepsilon(x,y),
 \end{aligned}$$

Some text after to test the below-display spacing.

```

\begin{equation*}
\begin{split}
f_{\{h,\varepsilon\}}(x,y)
&=\varepsilon\mathbf{E}_{\{x,y\}}\int_0^{t_\varepsilon}L_{\{x,y_\varepsilon(\varepsilon u)\}}\varphi(x)\,du\\
&= h\int L_{\{x,z\}}\varphi(x)\rho_x(dz)\\
&\quad +h\biggl[\frac{1}{t_\varepsilon}\biggl(\mathbf{E}_y\int_0^{t_\varepsilon}L_{\{x,y^x(s)\}}\varphi(x)\,ds\\
&\quad -t_\varepsilon\int L_{\{x,z\}}\varphi(x)\rho_x(dz)\biggr)\\
&\quad +\phantom{=}\frac{1}{t_\varepsilon}\biggl(\mathbf{E}_y\int_0^{t_\varepsilon}L_{\{x,y^x(s)\}}\varphi(x)\,ds\\
&\quad -\mathbf{E}_{\{x,y\}}\int_0^{t_\varepsilon}L_{\{x,y_\varepsilon(\varepsilon s)\}}\varphi(x)\,ds\biggr)\biggr] \\
&=h\mathbf{L}_x\varphi(x)+h\theta_\varepsilon(x,y),
\end{split}
\end{equation*}

```

If the option `centertags` is included in the options list of the `amsmath` package, the equation numbers for `split` environments will be centered vertically on the height of the `split`:

$$\begin{aligned}
 |I_2| &= \left| \int_0^T \psi(t) \left\{ u(a, t) - \int_{\gamma(t)}^a \frac{d\theta}{k(\theta, t)} \int_a^\theta c(\xi) u_t(\xi, t) d\xi \right\} dt \right| \\
 &\leq C_6 \left\| f \int_\Omega \left| \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right| \right\| \left\| |u| \overset{\circ}{\rightarrow} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\|.
 \end{aligned} \tag{65}$$

Some text after to test the below-display spacing.

Use of `split` within `align`:

$$\begin{aligned}
 |I_1| &= \left| \int_{\Omega} g R u \, d\Omega \right| \\
 &\leq C_3 \left[\int_{\Omega} \left(\int_a^x g(\xi, t) \, d\xi \right)^2 d\Omega \right]^{1/2} \\
 &\quad \times \left[\int_{\Omega} \left\{ u_x^2 + \frac{1}{k} \left(\int_a^x c u_t \, d\xi \right)^2 \right\} c \Omega \right]^{1/2} \\
 &\leq C_4 \left\| f \right\| \left\| \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right\| \left\| |u| \overset{\circ}{\rightarrow} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\|.
 \end{aligned} \tag{66}$$

$$\begin{aligned}
 |I_2| &= \left| \int_0^T \psi(t) \left\{ u(a, t) - \int_{\gamma(t)}^a \frac{d\theta}{k(\theta, t)} \int_a^{\theta} c(\xi) u_t(\xi, t) \, d\xi \right\} dt \right| \\
 &\leq C_6 \left\| f \right\| \left\| \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right\| \left\| |u| \overset{\circ}{\rightarrow} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\|.
 \end{aligned} \tag{67}$$

Some text after to test the below-display spacing.

```

\begin{align}
\begin{split}
&\&=\left\lvert\int_{\Omega} gRu\,,d\Omega\right\rvert\\
&\leq C_3\left[\int_{\Omega}\left(\int_a^x g(\xi,t)\,d\xi\right)^2d\Omega\right]^{1/2}\\
&\quad\times\left[\int_{\Omega}\left\{u_x^2+\frac{1}{k}\left(\int_a^x cu_t\,d\xi\right)^2\right\}c\Omega\right]^{1/2}\\
&\leq C_4\left\|f\right\|\left\|\tilde{S}_{a,-}^{-1,0}W_2(\Omega,\Gamma_l)\right\|\left\||u|\overset{\circ}{\rightarrow}W_2^{\tilde{A}}(\Omega;\Gamma_r,T)\right\|.
\end{split}
\end{align}
\label{eq:A}

\begin{split}
&\&=\left\lvert\int_0^T\psi(t)\left\{u(a,t)-\int_{\gamma(t)}^a\frac{d\theta}{k(\theta,t)}\int_a^{\theta}c(\xi)u_t(\xi,t)\,d\xi\right\}dt\right\rvert\\
&\leq C_6\left\|f\right\|\left\|\tilde{S}_{a,-}^{-1,0}W_2(\Omega,\Gamma_l)\right\|\left\||u|\overset{\circ}{\rightarrow}W_2^{\tilde{A}}(\Omega;\Gamma_r,T)\right\|.
\end{split}
\end{align}

```

Unnumbered align, with a number on the second split:

$$\begin{aligned}
 |I_1| &= \left| \int_{\Omega} g R u \, d\Omega \right| \\
 &\leq C_3 \left[\int_{\Omega} \left(\int_a^x g(\xi, t) \, d\xi \right)^2 d\Omega \right]^{1/2} \\
 &\quad \times \left[\int_{\Omega} \left\{ u_x^2 + \frac{1}{k} \left(\int_a^x c u_t \, d\xi \right)^2 \right\} c \Omega \right]^{1/2} \\
 &\leq C_4 \left\| f \left| \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right| \right\| \left\| |u| \xrightarrow{\circ} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\|. \\
 |I_2| &= \left| \int_0^T \psi(t) \left\{ u(a, t) - \int_{\gamma(t)}^a \frac{d\theta}{k(\theta, t)} \int_a^{\theta} c(\xi) u_t(\xi, t) \, d\xi \right\} dt \right| \\
 &\leq C_6 \left\| f \int_{\Omega} \left| \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right| \right\| \left\| |u| \xrightarrow{\circ} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\|.
 \end{aligned} \tag{67'}$$

Some text after to test the below-display spacing.

```

\begin{align*}
\begin{split}
&\abs{I_1} = \left\lvert \int_{\Omega} g R u \, d\Omega \right\rvert \\
&\leq C_3 \left[ \int_{\Omega} \left( \int_a^x g(\xi, t) \, d\xi \right)^2 d\Omega \right]^{1/2} \\
&\phantom{=} \times \left[ \int_{\Omega} \left\{ u_x^2 + \frac{1}{k} \left( \int_a^x c u_t \, d\xi \right)^2 \right\} c \Omega \right]^{1/2} \\
&\leq C_4 \left\lvert \left\lvert \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right\rvert \right\rvert \left\lvert |u| \xrightarrow{\circ} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\rvert.
\end{split} \\
\begin{split}
&\abs{I_2} = \left\lvert \int_0^T \psi(t) \left\{ u(a, t) - \int_{\gamma(t)}^a \frac{d\theta}{k(\theta, t)} \int_a^{\theta} c(\xi) u_t(\xi, t) \, d\xi \right\} dt \right\rvert \\
&\leq C_6 \left\lvert \int_{\Omega} \left\lvert \tilde{S}_{a,-}^{-1,0} W_2(\Omega, \Gamma_l) \right\rvert \right\rvert \left\lvert |u| \xrightarrow{\circ} W_2^{\tilde{A}}(\Omega; \Gamma_r, T) \right\rvert.
\end{split}
\end{align*}
\tag{67'}

```

A.2 Multline

Numbered version:

$$\begin{aligned} \int_a^b \left\{ \int_a^b [f(x)^2 g(y)^2 + f(y)^2 g(x)^2] - 2f(x)g(x)f(y)g(y) dx \right\} dy \\ = \int_a^b \left\{ g(y)^2 \int_a^b f^2 + f(y)^2 \int_a^b g^2 - 2f(y)g(y) \int_a^b fg \right\} dy \quad (68) \end{aligned}$$

To test the use of `\label` and `\ref`, we refer to the number of this equation here: (68).

```
\begin{multline}\label{eq:E}
\int_a^b\biggl\{\int_a^b[f(x)^2g(y)^2+f(y)^2g(x)^2]
-2f(x)g(x)f(y)g(y)\,dx\biggr\}\,dy \\\
=\int_a^b\biggl\{g(y)^2\int_a^bf^2+f(y)^2
\int_a^bg^2-2f(y)g(y)\int_a^b fg\biggr\}\,dy
\end{multline}
```

Unnumbered version:

$$\begin{aligned} \int_a^b \left\{ \int_a^b [f(x)^2 g(y)^2 + f(y)^2 g(x)^2] - 2f(x)g(x)f(y)g(y) dx \right\} dy \\ = \int_a^b \left\{ g(y)^2 \int_a^b f^2 + f(y)^2 \int_a^b g^2 - 2f(y)g(y) \int_a^b fg \right\} dy \end{aligned}$$

Some text after to test the below-display spacing.

```
\begin{multline*}
\int_a^b\biggl\{\int_a^b[f(x)^2g(y)^2+f(y)^2g(x)^2]
-2f(x)g(x)f(y)g(y)\,dx\biggr\}\,dy \\\
=\int_a^b\biggl\{g(y)^2\int_a^bf^2+f(y)^2
\int_a^bg^2-2f(y)g(y)\int_a^b fg\biggr\}\,dy
\end{multline*}
```

A.3 Gather

Numbered version with \notag on the second line:

$$D(a, r) \equiv \{z \in \mathbf{C}: |z - a| < r\}, \quad (69)$$

$$\text{seg}(a, r) \equiv \{z \in \mathbf{C}: \Im z = \Im a, |z - a| < r\},$$

$$c(e, \theta, r) \equiv \{(x, y) \in \mathbf{C}: |x - e| < y \tan \theta, 0 < y < r\}, \quad (70)$$

$$C(E, \theta, r) \equiv \bigcup_{e \in E} c(e, \theta, r). \quad (71)$$

```
\begin{gather}
D(a,r)\equiv\{z\in\mathbf{C}\colon \abs{z-a}<r\},\\
\text{seg}(a,r)\equiv\{z\in\mathbf{C}\colon \\
\Im z= \Im a, \abs{z-a}<r\},\notag\\
c(e,\theta,r)\equiv\{(x,y)\in\mathbf{C}\\
\colon \abs{x-e}<y\tan\theta, \ 0<y<r\},\\
C(E,\theta,r)\equiv\bigcup_{e\in E}c(e,\theta,r).
\end{gather}
```

Unnumbered version.

$$D(a, r) \equiv \{z \in \mathbf{C}: |z - a| < r\},$$

$$\text{seg}(a, r) \equiv \{z \in \mathbf{C}: \Im z = \Im a, |z - a| < r\},$$

$$c(e, \theta, r) \equiv \{(x, y) \in \mathbf{C}: |x - e| < y \tan \theta, 0 < y < r\},$$

$$C(E, \theta, r) \equiv \bigcup_{e \in E} c(e, \theta, r).$$

Some text after to test the below-display spacing.

```
\begin{gather*}
D(a,r)\equiv\{z\in\mathbf{C}\colon \abs{z-a}<r\},\\
\text{seg}(a,r)\equiv\{z\in\mathbf{C}\colon \\
\Im z= \Im a, \abs{z-a}<r\},\\
c(e,\theta,r)\equiv\{(x,y)\in\mathbf{C}\\
\colon \abs{x-e}<y\tan\theta, \ 0<y<r\},\\
C(E,\theta,r)\equiv\bigcup_{e\in E}c(e,\theta,r).
\end{gather*}
```

A.4 Align

Numbered version:

$$\gamma_x(t) = (\cos tu + \sin tx, v), \quad (72)$$

$$\gamma_y(t) = (u, \cos tv + \sin ty), \quad (73)$$

$$\gamma_z(t) = \left(\cos tu + \frac{\alpha}{\beta} \sin tv, -\frac{\beta}{\alpha} \sin tu + \cos tv \right). \quad (74)$$

Some text after to test the below-display spacing.

```
\begin{align}
\gamma_x(t)&=(\cos tu+\sin tx,v),\\
\gamma_y(t)&=(u,\cos tv+\sin ty),\\
\gamma_z(t)&=\left(\cos tu+\frac{\alpha}{\beta}\sin tv,
-\frac{\beta}{\alpha}\sin tu+\cos tv\right).
\end{align}
```

Unnumbered version:

$$\gamma_x(t) = (\cos tu + \sin tx, v),$$

$$\gamma_y(t) = (u, \cos tv + \sin ty),$$

$$\gamma_z(t) = \left(\cos tu + \frac{\alpha}{\beta} \sin tv, -\frac{\beta}{\alpha} \sin tu + \cos tv \right).$$

Some text after to test the below-display spacing.

```
\begin{align*}
\gamma_x(t)&=(\cos tu+\sin tx,v),\\
\gamma_y(t)&=(u,\cos tv+\sin ty),\\
\gamma_z(t)&=\left(\cos tu+\frac{\alpha}{\beta}\sin tv,
-\frac{\beta}{\alpha}\sin tu+\cos tv\right).
\end{align*}
```

A variation:

$$x = y \quad \text{by (84)} \quad (75)$$

$$x' = y' \quad \text{by (85)} \quad (76)$$

$$x + x' = y + y' \quad \text{by Axiom 1.} \quad (77)$$

Some text after to test the below-display spacing.

```
\begin{align}
x&=y \ \&\ \text{by (\ref{eq:C})}\\
x'&=y' \ \&\ \text{by (\ref{eq:D})}\\
x+x'&=y+y' \ \&\ \text{by Axiom 1.}
\end{align}
```


A.5 Align and split within gather

When using the `align` environment within the `gather` environment, one or the other, or both, should be unnumbered (using the `*` form); numbering both the outer and inner environment would cause a conflict.

Automatically numbered `gather` with `split` and `align*`:

$$\begin{aligned}\varphi(x, z) &= z - \gamma_{10}x - \gamma_{mn}x^m z^n \\ &= z - Mr^{-1}x - Mr^{-(m+n)}x^m z^n \\ \zeta^0 &= (\xi^0)^2, \\ \zeta^1 &= \xi^0 \xi^1, \\ \zeta^2 &= (\xi^1)^2,\end{aligned}\tag{78}$$

Here the `split` environment gets a number from the outer `gather` environment; numbers for individual lines of the `align*` are suppressed because of the star.

```
\begin{gather}
\begin{split} \varphi(x, z)
&= z - \gamma_{10}x - \gamma_{mn}x^m z^n \\
&= z - Mr^{-1}x - Mr^{-(m+n)}x^m z^n
\end{split} \\
\end{gather}
\begin{align*}
\zeta^0 &= (\xi^0)^2, \\
\zeta^1 &= \xi^0 \xi^1, \\
\zeta^2 &= (\xi^1)^2,
\end{align*}
\end{gather}
```

The `*`-ed form of `gather` with the non-`*`-ed form of `align`.

$$\begin{aligned}\varphi(x, z) &= z - \gamma_{10}x - \gamma_{mn}x^m z^n \\ &= z - Mr^{-1}x - Mr^{-(m+n)}x^m z^n \\ \zeta^0 &= (\xi^0)^2, \\ \zeta^1 &= \xi^0 \xi^1, \\ \zeta^2 &= (\xi^1)^2,\end{aligned}\tag{79}$$

(79)

(80)

(81)

Some text after to test the below-display spacing.

```
\begin{gather*}
\begin{split} \varphi(x, z)
&= z - \gamma_{10}x - \gamma_{mn}x^m z^n \\
&= z - Mr^{-1}x - Mr^{-(m+n)}x^m z^n
\end{split} \\
\end{gather*}
\begin{align}
\zeta^0 &= (\xi^0)^2, \\
\zeta^1 &= \xi^0 \xi^1, \\
\zeta^2 &= (\xi^1)^2,
\end{align}
\end{gather*}
```

A.6 Alignat

Numbered version:

$$V_i = v_i - q_i v_j, \quad X_i = x_i - q_i x_j, \quad U_i = u_i, \quad \text{for } i \neq j; \quad (82)$$

$$V_j = v_j, \quad X_j = x_j, \quad U_j u_j + \sum_{i \neq j} q_i u_i. \quad (83)$$

Some text after to test the below-display spacing.

```
\begin{alignat}{3}
V_i &= v_i - q_i v_j, & \quad \quad X_i &= x_i - q_i x_j, \\
& \quad \quad U_i &= u_i, \\
& \quad \quad \text{\textit{for } $i \neq j$}} \label{eq:B} \\
V_j &= v_j, & \quad \quad X_j &= x_j, \\
& \quad \quad U_j &= u_j + \sum_{i \neq j} q_i u_i. \\
\end{alignat}
```

Unnumbered version:

$$V_i = v_i - q_i v_j, \quad X_i = x_i - q_i x_j, \quad U_i = u_i, \quad \text{for } i \neq j;$$

$$V_j = v_j, \quad X_j = x_j, \quad U_j u_j + \sum_{i \neq j} q_i u_i.$$

Some text after to test the below-display spacing.

```
\begin{alignat*}{3}
V_i &= v_i - q_i v_j, & \quad \quad X_i &= x_i - q_i x_j, \\
& \quad \quad U_i &= u_i, \\
& \quad \quad \text{\textit{for } $i \neq j$}} \\
V_j &= v_j, & \quad \quad X_j &= x_j, \\
& \quad \quad U_j &= u_j + \sum_{i \neq j} q_i u_i. \\
\end{alignat*}
```

The most common use for `alignat` is for things like

$$x = y \quad \text{by (66)} \quad (84)$$

$$x' = y' \quad \text{by (82)} \quad (85)$$

$$x + x' = y + y' \quad \text{by Axiom 1.} \quad (86)$$

Some text after to test the below-display spacing.

```
\begin{alignat}{2}
x&=y && \quad \text{by (\ref{eq:A})}\label{eq:C}\\
x'&=y' && \quad \text{by (\ref{eq:B})}\label{eq:D}\\
x+x'&=y+y' && \quad \text{by Axiom 1.}
\end{alignat}
```

References

- [1] W. Diffie and E. Hellman, *New directions in cryptography*, IEEE Transactions on Information Theory **22** (1976), no. 5, 644–654.
- [2] D. H. Fremlin, *Cichon's diagram*, 1983/1984, presented at the Séminaire Initiation à l'Analyse, G. Choquet, M. Rogalski, J. Saint Raymond, at the Université Pierre et Marie Curie, Paris, 23e année.
- [3] I. P. Goulden and D. M. Jackson, *The enumeration of directed closed Euler trails and directed Hamiltonian circuits by Langrangian methods*, European J. Combin. **2** (1981), 131–212.
- [4] F. Harary and E. M. Palmer, *Graphical enumeration*, Academic Press, 1973.
- [5] R. Impagliazzo, L. Levin, and M. Luby, *Pseudo-random generation from one-way functions*, Proc. 21st STOC (1989), ACM, New York, pp. 12–24.
- [6] M. Kojima, S. Mizuno, and A. Yoshise, *A new continuation method for complementarity problems with uniform p -functions*, Tech. Report B-194, Tokyo Inst. of Technology, Tokyo, 1987, Dept. of Information Sciences.
- [7] ———, *A polynomial-time algorithm for a class of linear complementarity problems*, Tech. Report B-193, Tokyo Inst. of Technology, Tokyo, 1987, Dept. of Information Sciences.
- [8] C. J. Liu and Yutze Chow, *On operator and formal sum methods for graph enumeration problems*, SIAM J. Algorithms Discrete Methods **5** (1984), 384–438.
- [9] M. Marcus and H. Minc, *A survey of matrix theory and matrix inequalities*, Complementary Series in Math. **14** (1964), 21–48.
- [10] S. Mizuno, A. Yoshise, and T. Kikuchi, *Practical polynomial time algorithms for linear complementarity problems*, Tech. Report 13, Tokyo Inst. of Technology, Tokyo, April 1988, Dept. of Industrial Engineering and Management.
- [11] R. D. Monteiro and I. Adler, *Interior path following primal-dual algorithms, part II: Quadratic programming*, August 1987, Working paper, Dept. of Industrial Engineering and Operations Research.
- [12] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, N.J., 1970.
- [13] Y. Ye, *Interior algorithms for linear, quadratic and linearly constrained convex programming*, Ph.D. thesis, Stanford Univ., Palo Alto, Calif., July 1987, Dept. of Engineering–Economic Systems, unpublished.

Long title

L. First Author^{*1}, L. Second Author^{**1,2}, and L. Third Author^{***2}

¹ First address

² Second address

Received 15 November 2003, revised 30 November 2003, accepted 2 December 2003

Published online 3 December 2003

Key words List, of, comma, separated, keywords.

PACS 04A25

This is an example input file. Comparing it with the output it generates can show you how to produce a simple document of your own.

Copyright line will be provided by the publisher

1 Introduction

The class file w-art.cls represents an adaptation of the $\text{\LaTeX}$ 2 ϵ -standard class file article.cls and the $\mathcal{A}\mathcal{M}\mathcal{S}$ class file amsart.cls with the size option 10pt to the specific requirements of journal production at WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim. It can be used through the $\text{\LaTeX}$ -command

`\documentclass[<abbr>,fleqn, other options]{w-art}`

where <abbr> is an abbreviation of the journal name.

Table 1 Abbreviations for journal names.

adp	Ann. Phys. (Leipzig)	Annalen der Physik
anac	Appl. Num. Anal. Comp. Math.	Applied Numerical Analysis and Computational Mathematics
cnp	Contrib. Plasma Phys.	Contributions to Plasma Physics
mn	Math. Nachr.	Mathematische Nachrichten
mlq	Math. Log. Quart.	Mathematical Logic Quarterly
pop	Fortschr. Phys.	Fortschritte der Physik
pss	phys. stat. sol.	physica status solidi
pssa	phys. stat. sol. (a)	physica status solidi (a)
pssb	phys. stat. sol. (b)	physica status solidi (b)
zamm	ZAMM · Z. Angew. Math. Mech.	Zeitschrift für Angewandte Mathematik und Mechanik

One difference to the standard layout is the indentation by 3 cc or 4 cc of floats (figures and tables) and mathematical environments (`\[...]`, `equation`, `multline`, ...). To achieve this effect the new floats `vchtable` and `vchfigure` were added which are to be used in combination with `\vchcaption`. The standard `table`, `figure`, and `\caption` commands are nevertheless still working. So if there is the need to place a table or figure over the full width of the page these floats may still be used. In order to get short captions flushed left in contrast to the standard centered form the class loads internally the `caption2.sty` package by Harald Axel Sommerfeldt with the options `nooneline`, `small`, `bf`.

* Corresponding author: e-mail: x.y@xxx.yyy.zz, Phone: +00 999 999 999, Fax: +00 999 999 999

** Second author footnote.

*** Third author footnote.

2 Required packages

This class requires the standard L^AT_EX packages `calc`, `sidecap`, and `caption2` and the $\mathcal{A}\mathcal{M}\mathcal{S}$ -L^AT_EX packages.¹

2.1 New documentclass options

separatedheads (default): This gives the normal section, subsection and subsubsection headings with white space above and below the heading.

embeddedheads: With the `embeddedheads` option all section headings on all levels will be typeset as run-in headings without numbering like the standard L^AT_EX paragraph. If the numbering shall remain one has to set `\setcounter{secnumdepth}{3}` explicitly in the preamble of the document.

autolastpage (default): The pagenumber of the last page will automatically be determined by the class-file. In the `\pagespan{ }{ }` command only the first entry should be entered. If the second entry is also given it will be ignored. The pagenumber of the lastpage will be used in the running head of the first page. In order to get correct results the document has to be run through L^AT_EX at least twice.

noautolastpage: The value of the second argument of the `\pagespan{ }{ }` command will be printed as the last pagenumber.

referee: Prints the document with a larger amount of interline whitespace.

2.2 Floating objects – figures and tables

We have two different table environments: `table` and `vchtable`. The same holds true for figure: `figure` and `vchfigure`. The `vch`-types including their captions (`vchcaption`) are typically leftindented by an amount equal to the indentation of mathematical formulas.

For the caption layout the `caption2.sty` package is preloaded.

Additionally the `sidecap` package of the L^AT_EX-distribution will be loaded with the option “rightcaption” by the `w-art` class. This package defines the `SCfigure` and `SCTable` environment for figures and tables with captions on one side.

2.2.1 Tables

The L^AT_EX code for Table 2 is

```
\begin{table}[htb]
\caption{The caption inside a table environment.}
\label{tab:2}\renewcommand{\arraystretch}{1.5}
\begin{tabular}{lll} \hline
Description 1 & Description 2 & Description \\\hline
Row 1, Col 1 & Row 1, Col 2 & Row 1, Col 3 \\\hline
Row 2, Col 1 & Row 2, Col 2 & Row 2, Col 3 \\\hline
\end{tabular}
\end{table}
```

The L^AT_EX code for Table 3 (a `vchtable`) is

¹ If these packages are not part of your installation you may download them from the nearest CTAN server.

Table 2 The caption inside a table environment.

Description 1	Description 2	Description
Row 1, Col 1	Row 1, Col 2	Row 1, Col 3
Row 2, Col 1	Row 2, Col 2	Row 2, Col 3

```

\begin{vchtable}[htb]
\pchcaption{The caption inside a vchtable environment.}
\label{tab:3}\renewcommand{\arraystretch}{1.5}
\begin{tabular}{lll} \hline
Description 1 & Description 2 & Description \\ \hline
Row 1, Col 1 & Row 1, Col 2 & Row 1, Col 3 \\
Row 2, Col 1 & Row 2, Col 2 & Row 2, Col 3 \\ \hline
\end{tabular}
\end{vchtable}

```

Table 3 The caption inside a vchtable environment.

Description 1	Description 2	Description
Row 1, Col 1	Row 1, Col 2	Row 1, Col 3
Row 2, Col 1	Row 2, Col 2	Row 2, Col 3


2.2.2 Figures

The $\text{\LaTeX}$ code for Fig. 1 is

```

\begin{figure}[htb]
\includegraphics[width=\textwidth, height=2cm]{empty.eps}
\caption{The usual figure environment. It ...}
\label{fig:1}
\end{figure}

```

**Fig. 1** The usual figure environment. It may be used for figures spanning the whole page width.

The $\text{\LaTeX}$ code for Fig. 2 (a vchfigure) is

```

\begin{vchfigure}[htb]
\includegraphics[width=.5\textwidth]{empty.eps}
\pchcaption{A vchfigure environment with a vchcaption.
Figure and caption are leftindented.}
\label{fig:2}
\end{vchfigure}

```




Fig. 2 A vchfigure environment with a vchcaption. Figure and caption are leftindented.

The $\text{\LaTeX}$ code for Fig. 3 and 4 is

```
\begin{figure}[htb]
\begin{minipage}[t]{.45\textwidth}
\includegraphics[width=\textwidth]{empty.eps}
\caption{Two figures side by side with different numbers.}
\label{fig:3}
\end{minipage}
\hfil
\begin{minipage}[t]{.45\textwidth}
\includegraphics[width=\textwidth]{empty.eps}
\caption{This is the second picture.}
\label{fig:4}
\end{minipage}
\end{figure}
```




Fig. 3 Two figures side by side with different numbers.




Fig. 4 This is the second picture.

The $\text{\LaTeX}$ code for Fig. 5a and b is

```
\begin{figure}[htb]
```



```

\includegraphics[width=.45\textwidth]{empty.eps}~a)
\hfil
\includegraphics[width=.45\textwidth]{empty.eps}~b)
\caption{Two figures with one number. The figures
are referred to as a) and b).}
\label{fig:5}
\end{figure}

```




Fig. 5 Two figures with one number. The figures are referred to as a) and b).

2.2.3 SCfigure and STable environments

The SCfigure and STable environment may be used as provided and described in the documentation of the sidecap package.

So a typical SCfigure environment would look as follows:

```

\begin{SCfigure}[<relwidth>][<float>]
\includegraphics[<options>]{filename.eps}%
\caption{Caption of a SCfigure.}
\label{fig:x1}      % Give a unique label
\end{SCfigure}

```

where `relwidth` (optional) is the caption width relative to the width of the figure or table. A large value (e.g., 50) reserves the maximum width that is possible.

And `float` (optional) is like the floating position parameter of the original table/figure environments. Default is `[tb]`.

The alignment rules are:

- Figures and tables on top of a page should be top aligned with the caption.
- Figures and tables on bottom of a page should be bottom aligned with the caption.

The $\LaTeX$ code for Fig. 6 is

```

\begin{SCfigure}[4][htb]
\includegraphics[width=.3\textwidth]{empty.eps}%
\caption{Caption of a SCfigure figure. These captions
are always bottom aligned.}
\label{fig:6}
\end{SCfigure}

```




Fig. 6 Caption of a SCfigure figure. These captions are always bottom aligned.

In order to print a figure and a table side by side the `\setfloattype` command is introduced. The $\LaTeX$ code for Fig. 7 and Table 4 is

```
\begin{figure}[htb]
\begin{minipage}{.45\textwidth}
\includegraphics[width=\textwidth]{empty.eps}
\caption{Figure and table side by side. This is the picture.}
\label{fig:8}
\end{minipage}
\hfil
\begin{minipage}{.45\textwidth}
\setfloattype{table}
\caption{This is the table. ...}
\label{tab:4}
\renewcommand{\arraystretch}{1.5}
\begin{tabular}{lll}
...
\end{tabular}
\end{minipage}
\end{figure}
```




Fig. 7 Figure and table side by side. This is the picture.

Table 4 This is the table. Picture and table are both numbered independently.

Description 1	Description 2	Description
Row 1, Col 1	Row 1, Col 2	Row 1, Col 3
Row 2, Col 1	Row 2, Col 2	Row 2, Col 3

3 Test of math environments

Equations are always left-aligned. Therefore the option `fleqn` is used for the `documentclass` command by default. Note that `fleqn` does not work with unnumbered displayed equations written as $Ax = b$, so please use $\left[Ax=b \right]$ or an `equation*` or `gather*` environment instead.

By default the equations are consecutively numbered. This may be changed by putting the following command inside the preamble

$$\backslash numberwithin{equation}{section}$$

The latex math display environment $\left[\dots \right]$

$$\sum_{i=1}^{\infty} \frac{1}{i^2}$$

An equation environment:

$$\sum_{i=1}^{\infty} \frac{1}{i^2} \tag{1}$$

For more mathematical commands and environments please refer to the document `*-tma.tex` and the documentation of the $\mathcal{MS}$ classes.

3.1 Some predefined theorem like environments

Some predefined theorem like environments may be used by loading the package `w-thm.sty`. This package will load by itself the package `amsthm.sty`. So it will be easy to define new theorem- and definition-like environments. For further details refer to the documentation of the `amsthm.sty` package.

Table 5 Some predefined theorem like environments.

environment	caption	theoremstyle
thm, theorem	Theorem	theorem
prop, proposition	Proposition	theorem
lem, lemma	Lemma	theorem
cor, corollary	Corollary	theorem
axiom	Axiom	theorem
defs, defn, definition	Definition	definition
example	Example	definition
rem, remark	Remark	definition
notation	Notation	definition

Theorem 3.1 *This is a theorem.*

Theorem 3.2 *Another theorem.*

Proof. This is a proof. □

Definition 3.3 This is a definition.

Proposition 3.4 *This is a proposition.*

Lemma 3.5 *This is a lemma.*

Corollary 3.6 *This is a corollary.*

Example 3.7 This is an example.

Remark 3.8 This is a remark.

3.2 Definition of new theorem like environments

Because w-thm.sty uses amsthm.sty the definition of new theorem like environments will be done in the same manner as in the amsthm package. The definition of

```
\theoremstyle{plain}
\newtheorem{criterion}{Criterion}
\theoremstyle{definition}
\newtheorem{condition}[theorem]{Condition}
```

inside the preamble of the document will give the following environments.

Criterion 1 *This is a Criterion.*

Condition 3.9 This is a Condition.

If the name of a predefined environment has to be changed it can be done by e.g. typing

```
\renewcommand{\definitionname}{Definitions}
```

after the `\begin{document}` command.

Acknowledgements An acknowledgement may be placed at the end of the article.

The style of the following references should be used in all documents.

References

- [1] C. L. Pekeris, Phys. Rev. **112**, 1649 (1958); Phys. Rev. **126**, 1470 (1962).
- [2] R. C. Jones, J. Opt. Soc. Am. **45**, 799–808 (1955).
E. C. Doerner, J. Opt. Soc. Am. **52**, 669–672 (1962).
- [3] V. R. Velasco, Electronic properties of Fibonacci quasi-periodic heterostructures, phys. stat. sol. (b) **232**, 71–75 (1999).
- [4] H. Ibach, Electron Energy Loss Spectrometers: The Technology of High Performance, Springer Series in Optical Sciences Vol. 63 (Springer, Berlin, 1990).
- [5] D. E. Knuth, Seminumerical Algorithms, The Art of Computer Programming Vol. 2, second ed. (Addison-Wesley, Reading, MA, 1981).
- [6] B. B. Mandelbrot, The Fractal Geometry of Nature (W. H. Freeman & Co, San Francisco, 1988), chap. 5, pp. 217–219.
- [7] G. Casati and J. Ford (eds.), Stochastic Behavior in Classical and Quantum Hamiltonian Systems, Lecture Notes in Physics Vol. 93 (Springer, Berlin, 1979).
- [8] D. J. Thouless, in: The Quantum Hall Effect, edited by R. Prange and S. Girvin (Springer, New York, 1987), p. 101.
- [9] M. Xie, in: Proceedings of the 1995 Particle Accelerator Conference, Dallas, TX, 1995 (IEEE, Piscataway, 1996), p. 183.
- [10] S. Huan et al., in: Proceedings of the 20th International Conference on the Physics of Semiconductors, Thessaloniki 1990, edited by E. M. Anastassakis and J. D. Joannopoulos (World Scientific, Singapore, 1991), p. 1369.
- [11] S. P. Parkin, Ph.D. thesis, Stanford University (1988).
- [12] T. Terrific, New Sorting Algorithms, Tech. Rep., Fanstord University, Computer Science Department, Fanstord, CA (1977).

Greek symbols – w-greek.sty

α	<code>\alpha</code>	θ	<code>\theta</code>	o	<code>o</code>	τ	<code>\tau</code>
β	<code>\beta</code>	ϑ	<code>\vartheta</code>	π	<code>\pi</code>	υ	<code>\upsilon</code>
γ	<code>\gamma</code>	ι	<code>\iota</code>	ϖ	<code>\varpi</code>	ϕ	<code>\phi</code>
δ	<code>\delta</code>	κ	<code>\kappa</code>	ρ	<code>\rho</code>	φ	<code>\varphi</code>
ϵ	<code>\epsilon</code>	λ	<code>\lambda</code>	ϱ	<code>\varrho</code>	χ	<code>\chi</code>
ε	<code>\varepsilon</code>	μ	<code>\mu</code>	σ	<code>\sigma</code>	ψ	<code>\psi</code>
ζ	<code>\zeta</code>	ν	<code>\nu</code>	ς	<code>\varsigma</code>	ω	<code>\omega</code>
η	<code>\eta</code>	ξ	<code>\xi</code>				
Γ	<code>\itGamma</code>	Λ	<code>\itLambda</code>	Σ	<code>\itSigma</code>	Ψ	<code>\itPsi</code>
Δ	<code>\itDelta</code>	Ξ	<code>\itXi</code>	Υ	<code>\itUpsilon</code>	Ω	<code>\itOmega</code>
Θ	<code>\itTheta</code>	Π	<code>\itPi</code>	Φ	<code>\itPhi</code>		

Table 1: Slanted greek letters

α	<code>\upalpha</code>	θ	<code>\uptheta</code>	o	<code>\upo</code>	τ	<code>\uptau</code>
β	<code>\upbeta</code>	ϑ	<code>\upvartheta</code>	π	<code>\uppi</code>	υ	<code>\upupsilon</code>
γ	<code>\upgamma</code>	ι	<code>\upiota</code>	ϖ	<code>\upvarpi</code>	ϕ	<code>\upphi</code>
δ	<code>\updelta</code>	κ	<code>\upkappa</code>	ρ	<code>\uprho</code>	φ	<code>\upvarphi</code>
ϵ	<code>\upepsilon</code>	λ	<code>\uplambda</code>	ϱ	<code>\upvarrho</code>	χ	<code>\upchi</code>
ε	<code>\varepsilon</code>	μ	<code>\upmu</code>	σ	<code>\upsigma</code>	ψ	<code>\uppsi</code>
ζ	<code>\upzeta</code>	ν	<code>\upnu</code>	ς	<code>\upvarsigma</code>	ω	<code>\upomega</code>
η	<code>\upeta</code>	ξ	<code>\upxi</code>				
Γ	<code>\Gamma</code>	Λ	<code>\Lambda</code>	Σ	<code>\Sigma</code>	Ψ	<code>\Psi</code>
Δ	<code>\Delta</code>	Ξ	<code>\Xi</code>	Υ	<code>\Upsilon</code>	Ω	<code>\Omega</code>
Θ	<code>\Theta</code>	Π	<code>\Pi</code>	Φ	<code>\Phi</code>		

Table 2: Upright greek letters

α	<code>\bm{\alpha}</code>	θ	<code>\bm{\theta}</code>	o	<code>\bm{o}</code>	τ	<code>\bm{\tau}</code>
β	<code>\bm{\beta}</code>	ϑ	<code>\bm{\vartheta}</code>	π	<code>\bm{\pi}</code>	υ	<code>\bm{\upsilon}</code>
γ	<code>\bm{\gamma}</code>	ι	<code>\bm{\iota}</code>	ϖ	<code>\bm{\varpi}</code>	ϕ	<code>\bm{\phi}</code>
δ	<code>\bm{\delta}</code>	κ	<code>\bm{\kappa}</code>	ρ	<code>\bm{\rho}</code>	φ	<code>\bm{\varphi}</code>
ϵ	<code>\bm{\epsilon}</code>	λ	<code>\bm{\lambda}</code>	ϱ	<code>\bm{\varrho}</code>	χ	<code>\bm{\chi}</code>
ε	<code>\bm{\varepsilon}</code>	μ	<code>\bm{\mu}</code>	σ	<code>\bm{\sigma}</code>	ψ	<code>\bm{\psi}</code>
ζ	<code>\bm{\zeta}</code>	ν	<code>\bm{\nu}</code>	ς	<code>\bm{\varsigma}</code>	ω	<code>\bm{\omega}</code>
η	<code>\bm{\eta}</code>	ξ	<code>\bm{\xi}</code>				
Γ	<code>\bm{\Gamma}</code>	Λ	<code>\bm{\Lambda}</code>	Σ	<code>\bm{\Sigma}</code>	Ψ	<code>\bm{\Psi}</code>
Δ	<code>\bm{\Delta}</code>	Ξ	<code>\bm{\Xi}</code>	Υ	<code>\bm{\Upsilon}</code>	Ω	<code>\bm{\Omega}</code>
Θ	<code>\bm{\Theta}</code>	Π	<code>\bm{\Pi}</code>	Φ	<code>\bm{\Phi}</code>		

Table 3: Boldface variants of slanted greek letters

α	<code>\pmb{\upalpha}</code>	θ	<code>\pmb{\uptheta}</code>	o	<code>\pmb{\up{o}}</code>	τ	<code>\pmb{\uptau}</code>
β	<code>\pmb{\upbeta}</code>	ϑ	<code>\pmb{\upvartheta}</code>	π	<code>\pmb{\uppi}</code>	υ	<code>\pmb{\upupsilon}</code>
γ	<code>\pmb{\upgamma}</code>	ι	<code>\pmb{\upiota}</code>	ϖ	<code>\pmb{\upvarpi}</code>	ϕ	<code>\pmb{\upphi}</code>
δ	<code>\pmb{\updelta}</code>	κ	<code>\pmb{\upkappa}</code>	ρ	<code>\pmb{\uprho}</code>	φ	<code>\pmb{\upvarphi}</code>
ϵ	<code>\pmb{\upepsilon}</code>	λ	<code>\pmb{\uplambda}</code>	ϱ	<code>\pmb{\varrho}</code>	χ	<code>\pmb{\upchi}</code>
ε	<code>\pmb{\varepsilon}</code>	μ	<code>\pmb{\upmu}</code>	σ	<code>\pmb{\upsigma}</code>	ψ	<code>\pmb{\uppsi}</code>
ζ	<code>\pmb{\upzeta}</code>	ν	<code>\pmb{\upnu}</code>	ς	<code>\pmb{\upvarsigma}</code>	ω	<code>\pmb{\upomega}</code>
η	<code>\pmb{\upeta}</code>	ξ	<code>\pmb{\upxi}</code>				
Γ	<code>\bm{\Gamma}</code>	Λ	<code>\bm{\Lambda}</code>	Σ	<code>\bm{\Sigma}</code>	Ψ	<code>\bm{\Psi}</code>
Δ	<code>\bm{\Delta}</code>	Ξ	<code>\bm{\Xi}</code>	Υ	<code>\bm{\Upsilon}</code>	Ω	<code>\bm{\Omega}</code>
Θ	<code>\bm{\Theta}</code>	Π	<code>\bm{\Pi}</code>	Φ	<code>\bm{\Phi}</code>		

Table 4: Boldface variants of upright greek letters